

- HORNE, C. 1869. Notes on the Common Grey Hornbill of India (*Meniceros bicornis*). Proceedings of the Zoological Society of London. 1869:241–243.
- HUME, A. O. 1890. The nests and eggs of Indian birds. Second Edition. Volume III. R. H. Porter, London, United Kingdom.
- KANNAN, R. 1994. Ecology and the conservation of the Great Pied Hornbill (*Buceros bicornis*) in the Western Ghats of southern India. Dissertation. University of Arkansas, Fayetteville, USA.
- KANNAN, R. AND D. A. JAMES. 1997. Breeding biology of the Great Pied Hornbill (*Buceros bicornis*) in the Anaimalai Hills of southern India. Journal of the Bombay Natural History Society 94:451–465.
- KEMP, A. C. 1979. A review of hornbills: biology and radiation. The Living Bird 17:105–136.
- KEMP, A. C. 1995. The hornbills. Oxford University Press, London, United Kingdom.
- POONSWAD, P., A. TSUJI, AND C. NGAMPONGSAL. 1987. A comparative study on breeding biology of sympatric hornbill species (Bucerotidae) in Thailand with implication for breeding in captivity. Pages 250–315 in Proceedings of the Delacour/IFCB Symposium on Breeding Birds in Captivity. Los Angeles, California, USA.
- POULSEN, H. 1970. Nesting behaviour of the Black-casqued Hornbill *Ceratogymna atrata* and the Great Hornbill *Buceros bicornis*. Ornis Scandinavica 1:11–15.
- SANCHEZ, P. A. 1976. Properties and management of soils in the tropics. John Wiley and Sons, New York, USA.
- SPECTOR, W. S. (Editor). 1956. The handbook of biological data. W. B. Saunders Co., New York, USA.
- STOTT, K. 1951. A nesting record of hornbills in captivity. Aviculture Magazine 57:113–118.
- THORMAHLEN, M. P. AND S. Y. HEALY. 1990. Breeding Great Hornbill (*Buceros bicornis*) at Sacramento Zoo. Regional Conference Proceedings. American Association of Zoological Parks and Aquariums, Wheeling, West Virginia, USA.

The Wilson Journal of Ornithology 119(1):121–124, 2007

First Description of the Nest, Eggs, and Breeding Behavior of the Mérida Tapaculo (*Scytalopus meridanus*)

Karie L. Decker,^{1,2} Alina M. Niklison,¹ and Thomas E. Martin¹

ABSTRACT.—We provide the first description of the nest, eggs, and breeding behavior of the Mérida Tapaculo (*Scytalopus meridanus*). Data are from one pair in the moist cloud forest of Yacambu National Park, Venezuela during April–May 2004. Two nests, constructed by the same pair, were globular in structure and consisted of mossy material placed in a rock crevice of a muddy rock wall. The eggs were cream colored with an average mass of 4.19 g. Clutch sizes were one in the first nest and two in the second. The species showed bi-parental care in nest building and incubation. Nest attentiveness (percent time spent on the nest incubating) averaged $83.4 \pm 14\%$ (SD). Average on and off bouts were 33.24 and 6.34 min, respectively. Received 22 December 2005. Accepted 11 August 2006.

Breeding biology and life history traits of tropical birds remain poorly known. Nests have not been found nor described for many

species limiting our knowledge and understanding of tropical life histories. Of approximately 40 currently recognized species of *Scytalopus*, nests of ~12 species have been described (Sclater and Salvin 1879; Skutch 1972; Stiles 1979; Hilty and Brown 1986; Sick 1993; Krabbe and Schulenberg 1997, 2003; Christian 2001; Young and Zuchowski 2003; Greeney and Gelis 2005; Greeney and Rombough 2005). *Scytalopus* is found throughout the Andes from Central America to Tierra del Fuego Island (Fjeldså and Krabbe 1990, Krabbe and Schulenberg 2003). We present data on life history traits (nest description, clutch size, egg mass, nest building, egg laying, and incubation investment) of *Scytalopus meridanus* in Yacambu National Park, a wet cloud forest of north central Venezuela (09° 42' N, 69° 42' W; 1,900 m elevation).

Scytalopus are elusive birds, often only detectable by sound (Hilty et al. 2003, Krabbe and Schulenberg 2003). *Scytalopus meridanus*, a small 16.5-g bird, similar to a wren, is

¹ USGS, Montana Cooperative Wildlife Research Unit, University of Montana, Missoula, MT 59812, USA.

² Corresponding author; e-mail: karie.decker@umontana.edu

known to scuttle on the ground among thick brush and grasses (Hilty et al. 2003). We observed individuals within 30 m of a creek, foraging low to the ground with short, rapid movements. Their call is a series of 25–30 quick sharp monotone ‘wick’ notes, similar to that of the Northern Flicker (*Colaptes auratus*), repeated for 10–15 sec.

OBSERVATIONS

The first nest, found on 16 April 2004, was 0.8 m above a creek bed in an obscured crevice about 6.25 cm deep into a muddy, rock face. The globular nest was at the end of this crevice and was composed of moss, rootlets, and decomposing leaves. The inside diameter and height were 6.8 and 6.2 cm, respectively, while the outside diameter and height were 10.5 and 8.5 cm, respectively. Incubation had commenced prior to finding the nest and the adults were incubating a single, cream-colored egg of unknown age with a mass of 4.07 g. We monitored the nest for 5 days during the incubation period until it was depredated on 21 April, and videotaped it once for 5.75 hrs (17 Apr, 0705–1345 hrs EST) following Martin (2002). In the video, *S. meridanus* exhibited simultaneous incubation exchange, revealing that both male and female incubate. Nest attentiveness (percent time spent on the nest incubating) averaged 82.2%, while incubation bouts averaged 33.13 min ($n = 11$); the mean off bout length was 7.26 min ($n = 11$).

The second nest, a re-nest of the same pair, was located on 4 May 2004 approximately 15 m from the first nest. The nest was 1.2 m above the bottom of the creek bed, in a similarly concealed crevice 7 cm deep, and constructed from similar material. The nest contained one egg when found and, after checking the nest daily for eight days, another egg was laid. The day the second egg was laid (12 May), the eggs of the second nest weighed 4.38 and 4.12 g. We videotaped the nest three times: once during the laying period and twice during incubation, totaling nearly 19 hrs (7 May, 0741–1345; 15 May, 0732–1352; and 23 May, 0711–1346 hrs). The first video revealed that during the laying period, both adults continued to build the nest after one egg had been laid. Both parents repeatedly brought small mossy material to the nest. In addition, both adults appeared to intermittently incubate the

single egg, but had an average attentiveness of only 45.2%. This incubation activity occurred only during the cool early hours between 0741 and 0920 hrs. From 0921–1345 hrs, the parents visited the nest only briefly, bringing nesting material, but did not incubate. The two incubation videos revealed that nest attentiveness on the third day of incubation (15 May) and the eleventh day (23 May) was comparable to that of the first nest of unknown incubation age (percent time on = 82.6 and 84.8%, respectively). Mean on and off bout lengths during first and second incubation videos (37.20 min on, 7.34 min off; and 29.38 min on, 4.43 min off, respectively) also were similar to the first nest. We monitored the nest daily for 8 days before the second egg was laid (4–12 May), and for 13 days after (12–25 May), until depredated, at which point the eggs had not yet hatched. The incubation periods documented for other *Scytalopus* are 15–23 days (De Santo et al. 2002, Krabbe and Schulenberg 2003); the period that we monitored this nest is well within this range.

DISCUSSION

Life history traits have been described for only a few other species of *Scytalopus*. Egg color and shape were similar to most other *Scytalopus* described to date, as was clutch size (Stiles 1979, Whitney 1994, De Santo et al. 2002, Krabbe and Schulenberg 2003). Nest composition, placement, and shape were consistent with other tapaculos (Stiles 1979, Whitney 1994, De Santo et al. 2002, Krabbe and Schulenberg 2003, Young and Zuchowski 2003, Greeney and Gelis 2005, Greeney and Rombough 2005). Many tropical birds have been reported to lay eggs on alternating days, some even with three days separating egg laying (Skutch 1976). Unlike any passerine of which we are aware, *S. meridanus* laid a second egg nearly one week after the first. We do not know if this pattern is typical, but may result from the large investment in eggs. Few records report detailed information on parental investment. We observed fairly high nest attentiveness not atypical of shared incubators (Martin 2002; TEM, unpubl. data). *Scytalopus meridanus* is sexually monomorphic and we could not ascertain if parents contributed evenly in nest attentiveness. However, length

TABLE 1. Reproductive attributes (means) of *Scytalopus meridanus* in Venezuela, 2004 (this study) and four other congeners in Central and South America (Krabbe and Schulenberg 2003).

Species	Clutch size	Egg volume (cm ³)	Egg mass (g)	Adult mass (g)	Egg mass/body mass (%)
<i>S. meridanus</i> ^a	1–2	4.064	4.190	16.5	25.39
<i>S. speluncae</i>	2	2.897	2.987 ^b	13.75 ^c	21.72
<i>S. indigoticus</i>	2	3.006	3.099 ^b	14.80 ^c	20.94
<i>S. magellanicus</i>	2–3	3.179	3.278 ^b	11.00 ^c	29.80
<i>S. griseicollis</i>	2	2.531	2.609 ^b	17.97 ^c	14.52

^a Sample sizes for *S. meridanus* are: clutch size (2), egg mass/volume (3), and adult mass (1).

^b Egg mass calculated from measurements reported in Krabbe and Schulenberg (2003) (sample sizes not given) using the equation in Van Noordwijk et al. (1981).

^c Information from descriptions in Krabbe and Schulenberg (2003) (sample sizes not given).

of alternating bouts of different individuals appeared similar.

Scytalopus meridanus laid remarkably large eggs relative to the size of the 16.5-g adult (Martin et al. 2006), about 25% of its body weight. We could not locate egg mass records for other species of *Scytalopus*, but egg length and width measurements as well as adult mass were available for four species (Krabbe and Schulenberg 2003). We developed a relative coefficient (1.031) between mass and volume using egg length, width, and mass measurements from nine passerine species in Argentina (TEM, unpubl. data) to estimate egg mass. We inserted this coefficient to get the equation: $\text{mass} = (0.5 \times \text{length} \times \text{width}^2) \times 1.031$ (adapted from Van Noordwijk et al. 1981). All five *Scytalopus* species appear to have relatively high reproductive investment in eggs, given their body mass (Table 1). The allometric relationship across other species in Venezuela (Martin et al. 2006) predicts an average egg mass of approximately 2.6 g based on the 16.5-g body mass of adult *Scytalopus meridanus*. This predicted egg mass is much less than that observed and indicates that *Scytalopus* lays a large egg, even compared with other tropical species.

Scytalopus joins the ranks of other species in endemic tropical families by having small clutch size and shared incubation that yields relatively high nest attentiveness. What may be unusual is the interval between laying eggs and the size of the egg.

ACKNOWLEDGMENTS

We thank J. J. Fontaine, A. D. Chalfoun, and the rest of the students and researchers in our lab for advice and comments on this manuscript, as well as M.

J. Foguet and J. R. Lang for field assistance. We are also grateful to Carlos Rengifo and Carlos Bosque for editorial comments and valuable suggestions. This study was made possible by support under NSF grants DEB-9981527 and DEB-0543178 to T. E. Martin. Permit numbers are DM/00000455 from FONACIT and PA-INP-0072.003 from INPARQUES.

LITERATURE CITED

- CHRISTIAN, D. G. 2001. Nests and nesting behavior of some little known Panamanian birds. *Ornitologia Neotropical* 12:327–336.
- DE SANTO, T. L., M. F. WILLSON, K. E. SIEVING, AND J. J. ARMESTO. 2002. Nesting biology of tapaculos (Rhinocryptidae) in fragmented south-temperate rainforests of Chile. *Condor* 104:482–495.
- FJELDSÅ, J. AND N. KRABBE. 1990. Birds of the high Andes. Copenhagen Zoological Museum, University of Copenhagen and Svendborg, Copenhagen, Denmark.
- GREENEY, H. F. AND R. A. GELIS. 2005. The nest and nestlings of the Long-tailed Tapaculo (*Scytalopus micropterus*). *Ornitología Colombiana* 3:88–91.
- GREENEY, H. F. AND C. J. F. ROMBOUGH. 2005. First nest of the Chusquea Tapaculo (*Scytalopus parkeri*) in southern Ecuador. *Ornitologia Neotropical* 16:439–440.
- HILTY, S. L. AND W. L. BROWN. 1986. A guide to the birds of Columbia. Princeton University Press, Princeton, New Jersey, USA.
- HILTY, S. L., J. A. GWYNNE, G. TUDOR, AND R. MAYER DE SCHAUENSEE. 2003. Birds of Venezuela. Second Edition. Princeton University Press, Princeton, New Jersey, USA.
- KRABBE, N. AND T. S. SCHULENBERG. 1997. Species limits and natural history of *Scytalopus* tapaculos (Rhinocryptidae), with descriptions of the Ecuadorian taxa, including three new species. *Ornithological Monographs* 48:47–88.
- KRABBE, N. AND T. S. SCHULENBERG. 2003. Family Rhinocryptidae (tapaculos). Pages 748–783 in *Handbook of the birds of the world. Volume 8. Broadbills to tapaculos* (J. del Hoyo, A. Elliot, and D. Christie, Editors). Lynx Edicions, Barcelona, Spain.

- MARTIN, T. E. 2002. A new view of avian life-history evolution tested on an incubation paradox. *Proceedings of the Royal Society of London. Series B* 269:309–316.
- MARTIN, T. E., R. D. BASSAR, S. K. BASSAR, J. J. FONTAINE, P. LLOYD, H. MATHEWSON, A. M. NIKLISON, AND A. CHALFOUN. 2006. Life history and ecological correlates of geographic variation in egg and clutch mass among passerine species. *Evolution* 60:390–398.
- SCLATER, P. L. AND O. SALVIN. 1879. On the birds collected by the late Mr. T. K. Salmon in the state of Antioquia, United States of Columbia. *Proceedings of the Zoological Society of London 1879*: 486–550.
- SICK, H. 1993. *Birds in Brazil: a natural history*. Princeton University Press, Princeton, New Jersey, USA.
- SKUTCH, A. F. 1972. *Studies of tropical American birds*. Publications of the Nuttall Ornithological Club, Number 10. Cambridge, Massachusetts, USA.
- SKUTCH, A. F. 1976. *Parent birds and their young*. University of Texas Press, Austin, USA.
- STILES, E. W. 1979. Nest and eggs of the White-browed Tapaculo (*Scytalopus superciliaris*). *Condor* 81:208.
- WHITNEY, B. M. 1994. A new *Scytalopus* tapaculo (Rhinocryptidae) from Boliva, with notes on other Bolivian members of the genus and the *magellanicus* complex. *Wilson Bulletin* 106:585–614.
- VAN NOORDWIJK, A. J., L. P. C. KEIZER, J. H. VAN BALLEN, AND W. SCHRLOO. 1981. Genetic variation in egg dimensions in a population of Great Tit. *Genetica* 55:221–232.
- YOUNG, B. E. AND W. ZUCHOWSKI. 2003. First description of the nest of the Silvery-fronted Tapaculo (*Scytalopus argentifrons*). *Wilson Bulletin* 115: 91–93.

The Wilson Journal of Ornithology 119(1):124–126, 2007

An Interspecific Foraging Association Between Nearctic-Neotropical Migrant Passerines in Bolivia

Rosalind B. Renfrew¹

ABSTRACT.—I present the first published record of a foraging association between Nearctic-neotropical migrant bird species during the austral summer in South America. I observed Barn Swallows (*Hirundo rustica*) and Cliff Swallows (*Petrochelidon pyrrhonota*) in February 2005 repeatedly foraging on aerial insects flushed by flocks of Bobolinks (*Dolichonyx oryzivorus*) settling onto soybean plants (*Glycine max*). Additional observations would be needed to distinguish this behavior between an opportunistic association and a commensal relationship. *Received 25 November 2005. Accepted 28 July 2006.*

Foraging associations assumed to be commensal between passerine species have been described within mixed species flocks (e.g., Hino 1998) and between non-flocking species (e.g., Willis 1972, Maxson and Maxson 1981, Robbins 1981). Here, I report the first documentation of a foraging association between two flocking Nearctic-neotropical migrant species during the austral summer.

¹ Vermont Institute of Natural Science, 6565 Woodstock Rd., P. O. Box 1281, Quechee, VT 05059, USA; e-mail: rrenfrew@vinsweb.org

On 11 February 2005 from 0900 to 1000 hrs EST in San Juan, depto. Santa Cruz, Bolivia, ~100 km northwest of the city of Santa Cruz, I observed a flock of ~1,000 Bobolinks (*Dolichonyx oryzivorus*) foraging in soybean (*Glycine max*) fields. This large aggregation consisted of a series of smaller (30–400 individuals) flocks that moved across the field by landing in the soybeans for 15–60 sec, lifting to 1–3 m above the soybeans, flying 25–50 m, landing again, and repeating this pattern. After each landing, ~10–30 Barn Swallows (*Hirundo rustica*) and 0–5 Cliff Swallows (*Petrochelidon pyrrhonota*) captured aerial insects above the Bobolink flock, foraging 1–5 m above the soybean canopy for 5–10 sec. During two subsequent walking transects (400 m), perpendicular to and intersecting the flight path of the Bobolink flocks, I flushed Pyralid moths (*Omiodes indicata* Fabricius) with every step. The other insect species that flushed above the canopy, the adult stage of the velvetbean caterpillar (*Anticarsia gemmatalis* Hübner), was not abundant. No other insect species were observed flying

above the soybean canopy. The insect fauna on soybean leaves appeared to be species-poor and dominated by Pyralid moths, with species of Coleoptera and Homoptera occasionally noted.

The “following” behavior I observed is frequently used by swallows, which are known to follow mammals, birds, humans, and farm equipment that flush prey. In some cases, concentrations of animals serve as cues to swallows that indicate local concentrations of food resources (Brown and Brown 1995, 1999). Once a few individual swallows locate a food source, they may serve to attract more swallows, known as “local enhancement” (Brown 1988). My observation of swallows feeding on lepidopterans is unusual, as lepidopterans have been previously reported as comprising only a small fraction of the diet of Cliff and Barn swallows (Beal 1918). However, diet for these opportunistic feeders varies depending on availability of prey items (Brown and Brown 1995, 1999).

The association between the three species, if observed on a regular basis, could be an example of commensal insectivory, whereby the swallows increase their foraging efficiency, while Bobolink foraging efficiency is unaltered. Bobolinks are not known to forage on aerial insects and were not observed feeding above the soybean canopy. They were eating caterpillars on the soybean leaves (RBR, pers. obs.), indicating they were not competing with swallows for food resources. I did not observe interspecific aggression or other direct interactions between the Bobolinks and swallows, suggesting the swallows were not interfering with or otherwise altering Bobolink activity.

There are two ways that swallows might be increasing their foraging efficiency by following Bobolinks. However, I did not measure swallow foraging efficiency with, versus without, Bobolinks as “beaters” (Rand 1954). Bobolink flock activity appeared to invariably increase the number of prey flying per unit area (prey density), and swallows may have had a greater chance of locating prey, resulting in more capture attempts per foraging bout or per unit time (increased feeding rate, Brown 1988). Alternatively, when insects are forced to fly upon being flushed, the proportion of successful capture attempts per time

spent foraging may be higher (increased success rate, Kushlan 1978).

This association was not sustained over time. On seven subsequent dates over 2 weeks, I observed monospecific flocks of Bobolinks using the same soybean field in the absence of swallows. On one of these dates, Bobolink flocks were also observed in two soybean fields ~3 km away, but swallows were not seen over or adjacent to the fields. I also observed swallows foraging over soybean fields in the absence of Bobolinks.

The association reported may occur sporadically in South American soybean fields when Bobolink and swallow distributions overlap spatially and temporally. Conditions under which this phenomenon may occur in soybean fields must include (1) caterpillars consumed by Bobolinks are available (e.g., after emergence and prior to application of insecticides), (2) flying insects preferred by swallows are available, and (3) Bobolink flocks are sufficiently large to flush prey in quantities that enable swallows to increase their foraging efficiency. Swallows may opportunistically use a “following” strategy to increase their foraging efficiency in different habitats with different beater species in South America.

ACKNOWLEDGMENTS

I thank A. M. Saavedra for her priceless assistance in the field and the Museo de Historia Natural Noel Kempff Mercado for providing logistical and administrative support in Bolivia. Dr. D. W. Winkler and C. C. Rimmer provided helpful comments on the manuscript. I am grateful to Michael Sabourin and Dr. Alma Solis (Smithsonian) for identifying insects. I thank the anonymous referees for their suggestions and comments that improved this manuscript.

LITERATURE CITED

- BEAL, F. E. L. 1918. Food habits of the swallows, a family of valuable native birds. U.S. Department of Agriculture Bulletin 619.
- BROWN, C. R. 1988. Social foraging in Cliff Swallows: local enhancement, risk sensitivity, competition and the avoidance of predators. *Animal Behavior* 36:780–792.
- BROWN, C. R. AND M. B. BROWN. 1995. Cliff Swallow (*Petrochelidon pyrrhonota*). The birds of North America. Number 149.
- BROWN, C. R. AND M. B. BROWN. 1999. Barn Swallow (*Hirundo rustica*). The birds of North America. Number 452.
- HINO, T. 1998. Mutualistic and commensal organization of avian mixed-species foraging flocks in a

- forest of western Madagascar. *Journal of Avian Biology* 29:17–24.
- KUSHLAN, J. A. 1978. Commensalism in the Little Blue Heron. *Auk* 95:677–681.
- MAXSON, J. AND A. D. MAXSON. 1981. Commensal foraging between Hairy and Pileated woodpeckers. *Journal of Field Ornithology* 52:62–63.
- RAND, A. L. 1954. Social feeding behavior of birds. *Fieldiana-Zoology* 36:1–71.
- ROBBINS, M. B. 1981. Two cases of commensal feeding between passerines. *Wilson Bulletin* 93:391–392.
- WILLIS, E. O. 1972. The behavior of Spotted Antbirds. *Ornithological Monographs* 10.

The Wilson Journal of Ornithology 119(1):126–128, 2007

Interspecific Egg-dumping by a Violet-green Swallow in an Active Western Bluebird Nest

Danika Kleiber,^{1,2,6} Jenne Turner,^{1,3} Amber E. Budden,^{1,4} and Janis L. Dickinson^{1,5}

ABSTRACT.—We observed a Violet-green Swallow (*Tachycineta thalassina*) laying an egg in an active Western Bluebird (*Sialia mexicana*) nest. The Western Bluebird male and not the female, was aggressive to the Violet-green Swallow but the swallow remained to lay the egg. This is the first documented incidence of which we are aware involving altricial interspecific egg-laying during the nestling phase. We suggest the timing of this event was more consistent with incidental egg deposition, or egg-dumping, than brood parasitism or nest usurpation. *Received 13 December 2005. Accepted 24 July 2006.*

Observations of birds laying eggs in nests of other species are of interest because this behavior, when timed appropriately and directed at nests where parents fail to recognize foreign eggs and offspring, may be the starting point for evolution of interspecific brood parasitism. Other explanations for this behavior are nest usurpation and egg-dumping, which is the deposition of an egg into another species' nest without any specific adaptive function. We collected video footage of a Vi-

olet-green Swallow (*Tachycineta thalassina*) laying an egg in an active Western Bluebird (*Sialia mexicana*) nest containing nine-day-old nestlings at Hastings Natural History Reservation, Carmel Valley, California (36° 22' N, 121° 34' W). Nest boxes have been monitored at Hastings and nearby ranches as part of a long-term study of Western Bluebirds since 1983 (Dickinson et al. 1996). Western Bluebirds are the primary box-nesting species at this study site, but other secondary cavity nesters use nest boxes at lower frequencies, including Ash-throated Flycatchers (*Myiarchus cinerascens*), Oak Titmice (*Baeolophus inornatus*), House Wrens (*Troglodytes aedon*), Bewick's Wren (*Thryomanes bewickii*) and Violet-green Swallows (Table 1).

We conducted a study of parental feeding behaviors during spring 2005 using video to identify patterns of resource allocation within

TABLE 1. Frequency of nest box use by secondary cavity-nesting species at Hastings Natural History Reservation, Carmel Valley, California, from 1983 to 2004.

Species	Nests (n)	Total use (%)
Western Bluebird	1,860	69
Ash-throated Flycatcher	298	11
Oak Titmouse	204	8
House Wren	175	7
Violet-green Swallow	122	5
Bewick's Wren	2	<1
Total nesting attempts (at least one egg)	2,661	100

¹ Hastings Natural History Reservation, 38601 E. Carmel Valley Rd., Carmel Valley, CA 93924, USA.

² University of British Columbia, Department of Forest Sciences, Forest Sciences Centre, 2424 Main Mall, Vancouver, BC V6T 1Z4, Canada.

³ 1420 Walnut St., Ste. 650, Philadelphia, PA 19102, USA.

⁴ Department of Zoology, University of Toronto, 25 Harbord St., Toronto, ON M5S 3G5, Canada.

⁵ Cornell Laboratory of Ornithology, 159 Sapsucker Woods Rd., Ithaca, NY 14850, USA.

⁶ Corresponding author; e-mail: danika@ekit.com

the nest. The day before filming we replaced the natal nest box and nest material with a box of the same dimensions containing a Western Bluebird nest, but with a Plexiglas side to accommodate the camera. A video camera and small LED light were positioned within a sleeve connected to the nest box on the morning of filming, which began at approximately 0530 hrs PST. Tapes were set on short play and recorded for a period of 120 min. During the first recording an experimenter collected simultaneous observational data on parental activity from a blind approximately 30 m away. Following tape completion, a second tape was inserted to record an additional 120 min of nestling provisioning observations. No additional observational data on parental activity were collected from outside the nest.

OBSERVATIONS

We collected video data on 27 May 2005 from a Western Bluebird nest containing five nine-day-old nestlings and a single unhatched Western Bluebird egg. In this instance, the time stamp on the video unit failed, and observations were timed using the video counter and calibrated using the corresponding observational data. The video nest watch began at 0520:00 hrs. At 0711:53 hrs, a female Violet-green Swallow entered the nest box and sat in the far left corner at the top of the nest cup. At 0712:46, the Western Bluebird male entered the box and proceeded to peck at the Violet-green Swallow's head six times before leaving the nest box at 0713:03. The Violet-green Swallow responded to the pecking attack by tucking her head down and remaining still. After 5 sec, the female Western Bluebird fed a nestling from the box entrance. She appeared to ignore the Violet-green Swallow and entered the box only to turn around, leaving 11 sec after arrival. The adult male and female Western Bluebirds fed the nestlings twice more while the swallow was in the nest box, but did so without entering the nest. At 0721:41, the first video tape ended and the second was inserted. We estimated the time from the ending of the first tape and beginning of the second to be approximately 30 sec, and added that time to our calculations. The Western Bluebird adults did not enter the box again while the Violet-green Swallow female was present. At approximately 0733 hrs, the Vio-

let-green Swallow left the nest box, leaving a white egg on the outer rim of the nest cup where she had been sitting. During the afternoon the video nest box and nest were replaced with the natal nest box and nest; the Violet-green Swallow egg was also transferred. When we returned to the nest the following day the Violet-green Swallow egg was lying broken in the bottom of the nest. There was no second Violet-green Swallow egg.

DISCUSSION

Three hypotheses are possible explanations for our observations: nest usurpation, brood parasitism, and interspecific egg-dumping. Nest usurpation is a possible explanation given that Western Bluebirds and Violet-green Swallows are both secondary cavity nesters whose breeding times overlap (Brown et al. 1992, Guinan et al. 2000). However, in this instance, timing of egg-laying and the observed behavior of the Violet-green swallow was not consistent with nest usurpation. Violet-green Swallows in our population use nest boxes at a low frequency (Table 1), but are common and presumably use natural cavities as nesting sites. Previously described nest usurpation interactions between Violet-green Swallows and Western Bluebirds do not match the behavior we observed. In Arizona, Brawn (1990) observed large groups of Violet-green Swallows taking over a nest box defended by a Western Bluebird, but a single Violet-green Swallow, such as we observed, was not successful in displacing Western Bluebirds. Successful use of a Western Bluebird box by a single pair of Violet-green Swallows has been observed, but was preceded by interspecific provisioning of the bluebird's young (Eltzroth and Robinson 1984).

Brood parasitism can also be ruled out due to improper timing of laying. We found no record of either intra- or interspecific brood parasitism by Violet-green Swallows in the literature (Brown et al. 1992). The placement of the egg outside the nest cup would be inconsistent with brood parasitism behavior.

It is more likely the Violet-green Swallow laid the egg because she was physiologically committed to do so but was unable to lay in her original nest, a behavior known as "egg-dumping" (Wiens 1971). There are many documented cases of interspecific egg-dumping in

avian species not known for brood parasitic behavior (Bailey 1886, Holcomb 1967, Wiens 1971, Gustafson 1975, Cannell and Harrington 1984, Littlefield 1984, Carter 1987, Sealy 1989). Females may lay in another cavity because they are unable to approach their own cavity at the time of laying, due to the presence of a predator, nest usurpation, or other disturbance. Based on monitoring of nest boxes at Hastings Reservation, 11.2% of 98 Violet-green Swallow nests with at least one egg failed before hatching from 1983 through 2005. This small incidence of nest failure during laying and incubation is only partially attributable to predation, but indicates that infrequent nest predation or disturbance during laying could account for the behavior we observed. Although there are no data on conspecific nest usurpation in Violet-green Swallows, nest usurpation has been well documented in Tree Swallows (*Tachycineta bicolor*) (Leffelaar 1985).

ACKNOWLEDGMENTS

We thank Wendy Fair and Caitlin Stern for helpful comments and Cachagua General Store for gastronomic support. This research was funded by NSF grant IBN-009702 to JLD. Edited video footage may be viewed at: http://www.birds.cornell.edu/birdhouse/nestboxcam/movies.home/vgsw_in.webl courtesy of Cornell Laboratory of Ornithology's "The Birdhouse Network."

LITERATURE CITED

- BAILEY, H. B. 1886. The Brown Thrush laying in the nest of the Wood Thrush. *Auk* 4:78.
- BRAWN, J. D. 1990. Interspecific competition and social behavior in Violet-green Swallows. *Auk* 107:606–608.
- BROWN, C. R., A. M. KNOTT, AND E. J. DAMROSE. 1992. Violet-green Swallow (*Tachycineta thalassina*). The birds of North America. Number 14.
- CANNELL, P. F. AND B. A. HARRINGTON. 1984. Interspecific egg dumping by a Great Egret and Black-crowned Night Herons. *Auk* 101:889–891.
- CARTER, M. D. 1987. An incident of brood parasitism by the Verdin. *Wilson Bulletin* 99:136.
- DICKINSON, J. L., W. D. KOENIG, AND F. A. PITELKA. 1996. Fitness consequences of helping behavior in the Western Bluebird. *Behavioral Ecology* 7:168–177.
- ELTZROTH, E. K. AND S. R. ROBINSON. 1984. Violet-green Swallows help Western Bluebirds at the nest. *Journal of Field Ornithology* 55:259–261.
- GUINAN, J. A., P. A. GOWATY, AND E. K. ELTZROTH. 2000. Western Bluebird (*Sialia mexicana*). The birds of North America. Number 510.
- GUSTAFSON, J. R. 1975. A Sage Sparrow egg in a Black-throated Sparrow nest. *Auk* 92:805–806.
- HOLCOMB, L. C. 1967. Mourning Dove egg in nest of catbird and robin. *Wilson Bulletin* 79:450–451.
- LEFFELAAR, D. AND R. J. ROBERTSON. 1985. Nest usurpation and female competition for breeding opportunities by Tree Swallows. *Wilson Bulletin* 97:221–224.
- LITTLEFIELD, C. D. 1984. Sandhill Crane incubates a Canada Goose egg. *Wilson Bulletin* 96:719.
- SEALY, S. G. 1989. Incidental "egg dumping" by the House Wren in a Yellow Warbler nest. *Wilson Bulletin* 101:491–493.
- WIENS, J. A. 1971. "Egg-dumping" by the Grasshopper Sparrow in a Savannah Sparrow nest. *Auk* 88:185–186.

The Wilson Journal of Ornithology 119(1):128–129, 2007

Western Bluebird Captures a Western Fence Lizard

Caitlin A. Stern^{1,2}

ABSTRACT.—I observed a male Western Bluebird (*Sialia mexicana*) capture a western fence lizard (*Sceloporus occidentalis*), the first reported incidence of vertebrate-directed raptorial behavior in the Western Bluebird. There are no previously published reports of Western Bluebirds capturing vertebrate prey, although

there is one previous report of a Western Bluebird carrying an unidentified lizard in the manner of a prey item, and a few reports of predation on vertebrates by the congeneric Eastern Bluebird (*Sialia sialis*). Received 9 January 2006. Accepted 28 June 2006.

¹ Hastings Natural History Reservation, 38601 E. Carmel Valley Rd., Carmel Valley, CA 93924, USA.

² Current address: Department of Neurobiology and Behavior, Cornell University, Ithaca, NY 14853, USA; e-mail: cas245@cornell.edu

Western Bluebirds (*Sialia mexicana*) are primarily insectivorous during the breeding season, and rely upon berry crops during winter (Guinan et al. 2000). Individual Eastern Bluebirds (*Sialia sialis*), a congeneric species,

which is also primarily dependent upon insects and small fruits (Gowaty and Plissner 1998), have occasionally been observed capturing vertebrate prey, such as snakes (Flanigan 1971) and shrews (Pinkowski 1974). Beal (1915) reported the bones of lizards and tree frogs in the stomachs of Eastern Bluebirds, as did Bent (1949). Braman and Pogue (2005) found a dead, 8.3-cm-long flat-headed snake (*Tantilla gracilis*) in an Eastern Bluebird nest box, suggesting a past predation event. Of 217 Western Bluebird stomachs examined by Beal (1915), not one contained evidence of vertebrate prey. Herlugson's (1982) analysis of Western Bluebird stomach contents similarly yielded no evidence of vertebrate prey items, and adults were not observed delivering vertebrate prey to nestlings. Gaylord (1995) observed a male Western Bluebird carrying an unidentified lizard in his bill; however, she did not observe the capture or consumption of the lizard. There have been no previous reported observations of Western Bluebirds capturing vertebrate prey.

OBSERVATIONS

Western Bluebirds have been color-banded and their breeding biology monitored at Hastings Natural History Reservation, Carmel Valley, California, since 1983 (Dickinson et al. 1996). While censusing a winter group on 11 October 2005, I observed a first-winter male Western Bluebird catch a ~5-cm-long western fence lizard (*Sceloporus occidentalis*; Stebbins 1966) in his bill at 0928 hrs PST. The bluebird was perched on top of a wooden fence post, and the lizard was clinging to the side of the post ~7 cm below the bluebird. The bluebird left his perch, hovering briefly beside the post as he seized the lizard around its middle with his bill. This maneuver was executed quickly, and the lizard did not attempt to flee, although it arched its body after it was already held in the bluebird's bill. Returning to his perch on the same fence post, the bluebird twice beat the lizard's head against the post. This appeared to be an attempt to stun or kill the lizard. At 0930 hrs, the bluebird flew with the now immobile lizard still in his bill onto the property of a private ranch. I was not able to follow and ascertain whether the bluebird consumed the lizard because this ranch is closed to researchers.

DISCUSSION

Western Bluebirds commonly beat invertebrate prey against a perch before consumption, and both Flanigan (1971) and Pinkowski (1974) reported Eastern Bluebirds beating vertebrate prey items against perches before consuming them. The observed bluebird male was clearly treating the lizard as a prey item.

That vertebrate-directed raptorial behavior by Western Bluebirds has not been reported previously, despite intensive observation of Western Bluebird behavior, suggests that, similar to Eastern Bluebirds, Western Bluebirds rarely capture vertebrate prey. To my knowledge, capture of vertebrate prey by the third member of the genus *Sialia* (i.e., Mountain Bluebird, *Sialia currucoides*) has not been reported.

ACKNOWLEDGMENTS

I thank J. L. Dickinson, D. L. Kleiber, W. M. Fair, and three anonymous reviewers for providing valuable comments on this manuscript.

LITERATURE CITED

- BEAL, F. E. L. 1915. Food of the robins and bluebirds of the United States. U.S. Department of Agriculture Biological Survey Bulletin 171.
- BENT, A. C. 1949. Life histories of North American thrushes, kinglets, and their allies. U.S. National Museum Bulletin 196.
- BRAMAN, S. C. AND D. W. POGUE. 2005. Eastern Bluebird provisions nestlings with flat-headed snake. *Wilson Bulletin* 117:100–101.
- DICKINSON, J. L., W. D. KOENIG, AND F. A. PITELKA. 1996. Fitness consequences of helping behavior in the Western Bluebird. *Behavioral Ecology* 7:168–177.
- FLANIGAN, A. B. 1971. Predation on snakes by Eastern Bluebird and Brown Thrasher. *Wilson Bulletin* 83: 441.
- GAYLORD, S. 1995. Lizard cuisine. *Sialia* 17:65.
- GOWATY, P. A. AND J. H. PLISSNER. 1998. Eastern Bluebird (*Sialia sialis*). *The birds of North America*. Number 381.
- GUINAN, J. A., P. A. GOWATY, AND E. K. ELTZROTH. 2000. Western Bluebird (*Sialia mexicana*). *The birds of North America*. Number 510.
- HERLUGSON, C. J. 1982. Food of adult and nestling Western and Mountain bluebirds. *Murrelet* 63:59–65.
- PINKOWSKI, B. C. 1974. Predation on a shrew by an Eastern Bluebird. *Wilson Bulletin* 86:83.
- STEBBINS, R. C. 1966. A field guide to western reptiles and amphibians. Houghton Mifflin Company, Boston, Massachusetts, USA.