

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use TRULICITY safely and effectively. See full prescribing information for TRULICITY.

TRULICITY (dulaglutide) injection, for subcutaneous use
Initial U.S. Approval: 2014

WARNING: RISK OF THYROID C-CELL TUMORS

See full prescribing information for complete boxed warning.

- Dulaglutide causes thyroid C-cell tumors in rats. It is unknown whether TRULICITY causes thyroid C-cell tumors, including medullary thyroid carcinoma (MTC), in humans as the human relevance of dulaglutide-induced rodent thyroid C-cell tumors has not been determined (5.1, 13.1).
- TRULICITY is contraindicated in patients with a personal or family history of MTC and in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2). Counsel patients regarding the potential risk of MTC and symptoms of thyroid tumors (4, 5.1).

RECENT MAJOR CHANGES

Indications and Usage (1)	2/2020
Warnings and Precautions, Diabetic Retinopathy (5.7)	2/2020
Dosage and Administration (2.1)	9/2020

INDICATIONS AND USAGE

TRULICITY® is a glucagon-like peptide-1 (GLP-1) receptor agonist indicated:

- as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus.
- to reduce the risk of major adverse cardiovascular events in adults with type 2 diabetes mellitus who have established cardiovascular disease or multiple cardiovascular risk factors.

Limitations of Use:

- Has not been studied in patients with a history of pancreatitis. Consider other antidiabetic therapies in these patients (1, 5.2).
- Not for treatment of type 1 diabetes mellitus (1).
- Not recommended in patients with severe gastrointestinal disease, including severe gastroparesis (1, 5.6).

DOSAGE AND ADMINISTRATION

- Initiate at 0.75 mg subcutaneously once weekly. Increase the dose to 1.5 mg once weekly for additional glycemic control.
- If additional glycemic control is needed, increase the dose to 3 mg once weekly after at least 4 weeks on the 1.5 mg dose (2.1).
- If additional glycemic control is needed, increase to the maximum dose of 4.5 mg once weekly after at least 4 weeks on the 3 mg dose. (2.1).
- If a dose is missed, administer the missed dose as soon as possible if there are at least 3 days (72 hours) until the next scheduled dose (2.1).
- Administer once weekly at any time of day with or without food (2.2).
- Inject subcutaneously in the abdomen, thigh, or upper arm (2.2).

DOSAGE FORMS AND STRENGTHS

- Injection: 0.75 mg/0.5 mL solution in a single-dose pen (3)
- Injection: 1.5 mg/0.5 mL solution in a single-dose pen (3)
- Injection: 3 mg/0.5 mL solution in a single-dose pen (3)
- Injection: 4.5 mg/0.5 mL solution in a single-dose pen (3)

CONTRAINDICATIONS

- Patients with a personal or family history of medullary thyroid carcinoma or in patients with Multiple Endocrine Neoplasia syndrome type 2 (4, 5.1).
- Patients with a prior serious hypersensitivity reaction to TRULICITY or any of the product components (4, 5.4).

WARNINGS AND PRECAUTIONS

- Thyroid C-cell Tumors:** See Boxed Warning (5.1).
- Pancreatitis:** Has been reported in clinical trials. Discontinue promptly if pancreatitis is suspected. Do not restart if pancreatitis is confirmed (5.2).
- Hypoglycemia:** When TRULICITY is used with an insulin secretagogue or insulin, consider lowering the dose of the sulfonylurea or insulin to reduce the risk of hypoglycemia (5.3).
- Hypersensitivity Reactions:** Serious hypersensitivity reactions (e.g., anaphylactic reactions and angioedema) have occurred. Discontinue TRULICITY and promptly seek medical advice (5.4).
- Acute Kidney Injury:** Monitor renal function in patients with renal impairment reporting severe adverse gastrointestinal reactions (5.5).
- Severe Gastrointestinal Disease:** Use may be associated with gastrointestinal adverse reactions, sometimes severe. Has not been studied in patients with severe gastrointestinal disease and is not recommended in these patients (5.6).
- Diabetic Retinopathy Complications:** Have been reported in a cardiovascular outcomes trial. Monitor patients with a history of diabetic retinopathy (5.7).

ADVERSE REACTIONS

The most common adverse reactions, reported in ≥5% of patients treated with TRULICITY are: nausea, diarrhea, vomiting, abdominal pain, and decreased appetite (6.1).

To report SUSPECTED ADVERSE REACTIONS, contact Eli Lilly and Company at 1-800-LillyRx (1-800-545-5979) or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

TRULICITY delays gastric emptying and has the potential to reduce the rate of absorption of concomitantly administered oral medications (7.1, 12.3).

USE IN SPECIFIC POPULATIONS

- Pregnancy:** TRULICITY should be used during pregnancy only if the potential benefit justifies the potential risk to fetus (8.1).

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved Medication Guide.

Revised: 09/2020

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: RISK OF THYROID C-CELL TUMORS

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Dosage

2.2 Important Administration Instructions

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Risk of Thyroid C-cell Tumors

5.2 Pancreatitis

5.3 Hypoglycemia with Concomitant Use of Insulin Secretagogues or Insulin

5.4 Hypersensitivity Reactions

5.5 Acute Kidney Injury

5.6 Severe Gastrointestinal Disease

5.7 Diabetic Retinopathy Complications in Patients with a History of Diabetic Retinopathy

6 ADVERSE REACTIONS

6.1 Clinical Studies Experience

6.2 Immunogenicity

6.3 Postmarketing Experience

7 DRUG INTERACTIONS

7.1 Oral Medications

7.2 Concomitant Use with an Insulin Secretagogue (e.g., Sulfonylurea) or with Insulin

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment
- 8.8 Gastroparesis

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, and Impairment of Fertility
- 13.2 Animal Toxicology and/or Pharmacology

14 CLINICAL STUDIES

- 14.1 Glycemic Control Trials in Adults with Type 2 Diabetes Mellitus
- 14.2 Cardiovascular Outcomes Trial in Adults with Type 2 Diabetes Mellitus and Cardiovascular Disease or Multiple Cardiovascular Risk Factors

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Storage and Handling

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: RISK OF THYROID C-CELL TUMORS

- In male and female rats, dulaglutide causes a dose-related and treatment-duration-dependent increase in the incidence of thyroid C-cell tumors (adenomas and carcinomas) after lifetime exposure. It is unknown whether TRULICITY causes thyroid C-cell tumors, including medullary thyroid carcinoma (MTC), in humans as human relevance of dulaglutide-induced rodent thyroid C-cell tumors has not been determined [see *Warnings and Precautions (5.1)*, and *Nonclinical Toxicology (13.1)*].
- TRULICITY is contraindicated in patients with a personal or family history of MTC and in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2). Counsel patients regarding the potential risk of MTC with use of TRULICITY and inform them of symptoms of thyroid tumors (e.g., mass in the neck, dysphagia, dyspnea, persistent hoarseness). Routine monitoring of serum calcitonin or using thyroid ultrasound is of uncertain value for early detection of MTC in patients treated with TRULICITY [see *Contraindications (4)* and *Warnings and Precautions (5.1)*].

1 INDICATIONS AND USAGE

TRULICITY® is indicated

- as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus.
- to reduce the risk of major adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) in adults with type 2 diabetes mellitus who have established cardiovascular disease or multiple cardiovascular risk factors.

Limitations of Use

- TRULICITY has not been studied in patients with a history of pancreatitis [see *Warnings and Precautions (5.2)*]. Consider other antidiabetic therapies in patients with a history of pancreatitis.
- TRULICITY should not be used in patients with type 1 diabetes mellitus.
- TRULICITY has not been studied in patients with severe gastrointestinal disease, including severe gastroparesis and is therefore not recommended in these patients [see *Warnings and Precautions (5.6)*].

2 DOSAGE AND ADMINISTRATION

2.1 Dosage

- The recommended initiating dose of TRULICITY is 0.75 mg injected subcutaneously once weekly.
- Increase the dose to 1.5 mg once weekly for additional glycemic control.
- If additional glycemic control is needed, increase the dose to 3 mg once weekly after at least 4 weeks on the 1.5 mg dose.
- If additional glycemic control is needed, increase the dose to the maximum dose of 4.5 mg once weekly after at least 4 weeks on the 3 mg dose.
- If a dose is missed, instruct patients to administer as soon as possible if there are at least 3 days (72 hours) until the next scheduled dose. If less than 3 days remain before the next scheduled dose, skip the missed dose and administer the next dose on the regularly scheduled day. In each case, patients can then resume their regular once weekly dosing schedule.
- The day of weekly administration can be changed if necessary, as long as the last dose was administered 3 or more days before.

2.2 Important Administration Instructions

- Administer TRULICITY once weekly, any time of day, with or without food.

- Inject TRULICITY subcutaneously in the abdomen, thigh, or upper arm.
- Rotate injection sites with each dose.
- Inspect TRULICITY visually before use. It should appear clear and colorless. Do not use TRULICITY if particulate matter or coloration is seen.
- When using TRULICITY with insulin, administer as separate injections and never mix. It is acceptable to inject TRULICITY and insulin in the same body region, but the injections should not be adjacent to each other.

3 DOSAGE FORMS AND STRENGTHS

Injection: TRULICITY is a clear and colorless solution available as:

- 0.75 mg/0.5 mL solution in a single-dose pen
- 1.5 mg/0.5 mL solution in a single-dose pen
- 3 mg/0.5 mL solution in a single-dose pen
- 4.5 mg/0.5 mL solution in a single-dose pen

4 CONTRAINDICATIONS

TRULICITY is contraindicated in patients with:

- Personal or family history of medullary thyroid carcinoma (MTC) or in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2) [see *Warnings and Precautions* (5.1)].
- Prior serious hypersensitivity reaction to dulaglutide or to any of the product components. Serious hypersensitivity reactions including anaphylactic reactions and angioedema have been reported with TRULICITY [see *Warnings and Precautions* (5.4)].

5 WARNINGS AND PRECAUTIONS

5.1 Risk of Thyroid C-cell Tumors

In male and female rats, dulaglutide causes a dose-related and treatment-duration-dependent increase in the incidence of thyroid C-cell tumors (adenomas and carcinomas) after lifetime exposure [see *Nonclinical Toxicology* (13.1)]. Glucagon-like peptide-1 (GLP-1) receptor agonists have induced thyroid C-cell adenomas and carcinomas in mice and rats at clinically relevant exposures. It is unknown whether TRULICITY will cause thyroid C-cell tumors, including medullary thyroid carcinoma (MTC), in humans, as the human relevance of dulaglutide-induced rodent thyroid C-cell tumors has not been determined.

One case of MTC was reported in a patient treated with TRULICITY in a clinical study. This patient had pretreatment calcitonin levels approximately 8 times the upper limit of normal (ULN). An additional case of C-cell hyperplasia with elevated calcitonin levels following treatment was reported in the cardiovascular outcomes trial (REWIND). Cases of MTC in patients treated with liraglutide, another GLP-1 receptor agonist, have been reported in the postmarketing period; the data in these reports are insufficient to establish or exclude a causal relationship between MTC and GLP-1 receptor agonist use in humans.

TRULICITY is contraindicated in patients with a personal or family history of MTC or in patients with MEN 2. Counsel patients regarding the potential risk for MTC with the use of TRULICITY and inform them of symptoms of thyroid tumors (e.g. a mass in the neck, dysphagia, dyspnea, persistent hoarseness).

Routine monitoring of serum calcitonin or using thyroid ultrasound is of uncertain value for early detection of MTC in patients treated with TRULICITY. Such monitoring may increase the risk of unnecessary procedures, due to the low test specificity for serum calcitonin and a high background incidence of thyroid disease. Significantly elevated serum calcitonin values may indicate MTC and patients with MTC usually have calcitonin values >50 ng/L. If serum calcitonin is measured and found to be elevated, the patient should be further evaluated. Patients with thyroid nodules noted on physical examination or neck imaging should also be further evaluated.

5.2 Pancreatitis

In a pooled analysis from the original registration studies, 12 (3.4 cases per 1000 patient years) pancreatitis-related adverse reactions were reported in patients exposed to TRULICITY versus 3 in non-incretin comparators (2.7 cases per 1000 patient years). An analysis of adjudicated events revealed 5 cases of confirmed pancreatitis in patients exposed to TRULICITY (1.4 cases per 1000 patient years) versus 1 case in non-incretin comparators (0.88 cases per 1000 patient years).

Based on an analysis of adjudicated events in a clinical study evaluating Trulicity 1.5 mg, 3 mg, or 4.5 mg once weekly, pancreatitis occurred in 1 patient exposed to TRULICITY 1.5 mg (0.2%), in 2 patients exposed to TRULICITY 3 mg (0.3%), and 3 patients exposed to TRULICITY 4.5 mg (0.5%).

After initiation of TRULICITY, observe patients carefully for signs and symptoms of pancreatitis, including persistent severe abdominal pain, sometimes radiating to the back, which may or may not be accompanied by vomiting. If pancreatitis is suspected, promptly discontinue TRULICITY. If pancreatitis is confirmed, TRULICITY should not be

restarted. TRULICITY has not been evaluated in patients with a prior history of pancreatitis. Consider other antidiabetic therapies in patients with a history of pancreatitis.

5.3 Hypoglycemia with Concomitant Use of Insulin Secretagogues or Insulin

The risk of hypoglycemia is increased when TRULICITY is used in combination with insulin secretagogues (e.g., sulfonylureas) or insulin. Patients may require a lower dose of sulfonylurea or insulin to reduce the risk of hypoglycemia in this setting [see *Adverse Reactions (6.1)*, *Drug Interactions (7.2)*].

5.4 Hypersensitivity Reactions

There have been postmarketing reports of serious hypersensitivity reactions including anaphylactic reactions and angioedema in patients treated with TRULICITY [see *Adverse Reactions (6.3)*]. If a hypersensitivity reaction occurs, discontinue TRULICITY; treat promptly per standard of care, and monitor until signs and symptoms resolve. Do not use in patients with a previous hypersensitivity reaction to TRULICITY [see *Contraindications (4)*].

Anaphylaxis and angioedema have been reported with other GLP-1 receptor agonists. Use caution in a patient with a history of angioedema or anaphylaxis with another GLP-1 receptor agonist because it is unknown whether such patients will be predisposed to anaphylaxis with TRULICITY.

5.5 Acute Kidney Injury

In patients treated with GLP-1 receptor agonists, including TRULICITY, there have been postmarketing reports of acute renal failure and worsening of chronic renal failure, which may sometimes require hemodialysis. Some of these events were reported in patients without known underlying renal disease. A majority of reported events occurred in patients who had experienced nausea, vomiting, diarrhea, or dehydration. Because these reactions may worsen renal function, use caution when initiating or escalating doses of TRULICITY in patients with renal impairment. Monitor renal function in patients with renal impairment reporting severe adverse gastrointestinal reactions [see *Use in Specific Populations (8.6)*].

5.6 Severe Gastrointestinal Disease

Use of TRULICITY may be associated with gastrointestinal adverse reactions, sometimes severe [see *Adverse Reactions (6.1)*]. TRULICITY has not been studied in patients with severe gastrointestinal disease, including severe gastroparesis, and is therefore not recommended in these patients.

5.7 Diabetic Retinopathy Complications in Patients with a History of Diabetic Retinopathy

In a cardiovascular outcomes trial with a median follow up of 5.4 years involving patients with type 2 diabetes with established cardiovascular disease or multiple cardiovascular risk factors, diabetic retinopathy complications occurred in patients treated with TRULICITY 1.5 mg (1.9%) and placebo (1.5%). These events were prospectively ascertained as a secondary composite endpoint. The proportion of patients with diabetic retinopathy complications was larger among patients with a history of diabetic retinopathy at baseline (TRULICITY 8.5%, placebo 6.2%) than among patients without a known history of diabetic retinopathy (TRULICITY 1%, placebo 1%).

Rapid improvement in glucose control has been associated with a temporary worsening of diabetic retinopathy. Patients with a history of diabetic retinopathy should be monitored for progression of diabetic retinopathy.

6 ADVERSE REACTIONS

The following serious reactions are described below or elsewhere in the prescribing information:

- Risk of Thyroid C-cell Tumors [see *Warnings and Precautions (5.1)*]
- Pancreatitis [see *Warnings and Precautions (5.2)*]
- Hypoglycemia with Concomitant Use of Insulin Secretagogues or Insulin [see *Warnings and Precautions (5.3)*]
- Hypersensitivity Reactions [see *Warnings and Precautions (5.4)*]
- Acute Kidney Injury [see *Warnings and Precautions (5.5)*]
- Severe Gastrointestinal Disease [see *Warnings and Precautions (5.6)*]
- Diabetic Retinopathy Complications in Patients with a History of Diabetic Retinopathy [see *Warnings and Precautions (5.7)*]

6.1 Clinical Studies Experience

Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice.

Pool of Placebo-controlled Trials for TRULICITY 0.75 mg and 1.5 mg Doses

The data in Table 1 are derived from a pool of placebo-controlled trials and include 1670 patients exposed to TRULICITY with a mean duration of exposure of 23.8 weeks [see *Clinical Studies (14)*]. The mean age of patients was 56 years, 1% were 75 years or older and 53% were male. The population was 69% White, 7% Black or African American, 13% Asian; 30% were of Hispanic or Latino ethnicity. At baseline, the population had diabetes for an average of 8 years, a

mean HbA1c of 8.0%, and 2.5% of the population reported retinopathy. Baseline estimated renal function was normal or mildly impaired (eGFR \geq 60 mL/min/1.73 m²) in 96%.

Table 1 shows adverse reactions, excluding hypoglycemia, occurring in \geq 5% of TRULICITY treated patients and more commonly than placebo in a pool of placebo-controlled trials.

Table 1: Adverse Reactions in Pool of Placebo-Controlled Trials Occurring in \geq 5% of TRULICITY-Treated Patients

Adverse Reaction	Placebo (N=568) %	TRULICITY 0.75 mg (N=836) %	TRULICITY 1.5 mg (N=834) %
Nausea	5.3	12.4	21.1
Diarrhea ^a	6.7	8.9	12.6
Vomiting ^b	2.3	6.0	12.7
Abdominal Pain ^c	4.9	6.5	9.4
Decreased Appetite	1.6	4.9	8.6
Dyspepsia	2.3	4.1	5.8
Fatigue ^d	2.6	4.2	5.6

^a Includes diarrhea, fecal volume increased, frequent bowel movements.

^b Includes retching, vomiting, vomiting projectile.

^c Includes abdominal discomfort, abdominal pain, abdominal pain lower, abdominal pain upper, abdominal tenderness, gastrointestinal pain.

^d Includes fatigue, asthenia, malaise.

Note: Percentages reflect the number of patients that reported at least 1 treatment-emergent occurrence of the adverse reaction.

Gastrointestinal Adverse Reactions

In the pool of placebo-controlled trials, gastrointestinal adverse reactions occurred more frequently among patients receiving TRULICITY than placebo (placebo 21.3%, 0.75 mg 31.6%, 1.5 mg 41.0%). More patients receiving TRULICITY 0.75 mg (1.3%) and TRULICITY 1.5 mg (3.5%) discontinued treatment due to gastrointestinal adverse reactions than patients receiving placebo (0.2%). Investigators graded the severity of gastrointestinal adverse reactions occurring on 0.75 mg and 1.5 mg of TRULICITY as “mild” in 58% and 48% of cases, respectively, “moderate” in 35% and 42% of cases, respectively, or “severe” in 7% and 11% of cases, respectively.

The following adverse reactions were reported more frequently in TRULICITY-treated patients than placebo (frequencies listed, respectively, as: placebo; 0.75 mg; 1.5 mg): constipation (0.7%, 3.9%, 3.7%), flatulence (1.4%, 1.4%, 3.4%), abdominal distension (0.7%, 2.9%, 2.3%), gastroesophageal reflux disease (0.5%, 1.7%, 2.0%), and eructation (0.2%, 0.6%, 1.6%).

TRULICITY 3 mg and 4.5 mg Doses

Table 2 shows adverse reactions occurring \geq 5% in any of the treatment groups through 36 weeks in a clinical trial with 1842 patients treated with Trulicity 1.5 mg, 3 mg, or 4.5 mg once weekly as an add-on to metformin. The adverse reaction profile is consistent with previous clinical trials.

Table 2: Adverse Reactions Occurring in \geq 5% of Patients Receiving TRULICITY 1.5 mg, 3 mg, or 4.5 mg in a Clinical Trial through 36 Weeks^a

Adverse Reaction	TRULICITY 1.5 mg (N=612) %	TRULICITY 3 mg (N=616) %	TRULICITY 4.5 mg (N=614) %
Nausea	13.4	15.6	16.4
Diarrhea	7.0	11.4	10.7
Vomiting	5.6	8.3	9.3
Dyspepsia	2.8	5.0	2.6

^a Percentages reflect the number of patients that reported at least 1 treatment-emergent occurrence of the adverse reaction.

Other Adverse Reactions

Hypoglycemia

Table 3 summarizes the incidence of hypoglycemia in the placebo-controlled clinical studies: episodes with a glucose level $<$ 54 mg/dL with or without symptoms, and severe hypoglycemia, defined as an episode requiring the assistance of another person to actively administer carbohydrate, glucagon, or other resuscitative actions.

Table 3: Incidence (%) of Hypoglycemia in Placebo-Controlled Trials

	Placebo	TRULICITY 0.75 mg	TRULICITY 1.5 mg
Add-on to Metformin			
(26 weeks)	N=177	N=302	N=304
Hypoglycemia with a glucose level <54 mg/dL	0	0.3	0.7
Severe hypoglycemia	0	0	0
Add-on to Metformin + Pioglitazone			
(26 weeks)	N=141	N=280	N=279
Hypoglycemia with a glucose level <54 mg/dL	1.4	2.1	0
Severe hypoglycemia	0	0	0
Add-on to Glimepiride			
(24 weeks)	N=60	-	N=239
Hypoglycemia with a glucose level <54 mg/dL	0	-	3.3
Severe hypoglycemia	0	-	0
In Combination with Insulin Glargine ± Metformin			
(28 weeks)	N=150	-	N=150
Hypoglycemia with a glucose level <54 mg/dL	9.3	-	14.7
Severe hypoglycemia	0	-	0.7
Add-on to SGLT2i ± Metformin			
(24 weeks)	N=140	N=141	N=142
Hypoglycemia with a glucose level <54 mg/dL	0.7	0.7	0.7
Severe hypoglycemia	0	0.7	0

Hypoglycemia was more frequent when TRULICITY was used in combination with a sulfonylurea or insulin than when used with non-secretagogues. In a 78-week clinical trial, hypoglycemia (glucose level <54 mg/dL) occurred in 20% and 21% of patients when TRULICITY 0.75 mg and 1.5 mg, respectively, were co-administered with a sulfonylurea. Severe hypoglycemia occurred in 0% and 0.7% of patients when TRULICITY 0.75 mg and 1.5 mg, respectively, were co-administered with a sulfonylurea. In a 52-week clinical trial, hypoglycemia (glucose level <54 mg/dL) occurred in 77% and 69% of patients when TRULICITY 0.75 mg and 1.5 mg, respectively, were co-administered with prandial insulin. Severe hypoglycemia occurred in 2.7% and 3.4% of patients when TRULICITY 0.75 mg and 1.5 mg, respectively, were co-administered with prandial insulin. Refer to Table 3 for the incidence of hypoglycemia in patients treated in combination with basal insulin glargine.

In the clinical trial with patients on TRULICITY 1.5 mg, TRULICITY 3 mg, or TRULICITY 4.5 mg once weekly, as add-on to metformin, incidences of hypoglycemia (glucose level <54 mg/dL) through 36 weeks were 1.1%, 0.3%, and 1.1%, respectively, and incidences of severe hypoglycemia were 0.2%, 0%, and 0.2%, respectively.

Cholelithiasis and Cholecystitis

In a cardiovascular outcomes trial with a median follow up of 5.4 years, cholelithiasis occurred at a rate of 0.62/100 patient-years in TRULICITY-treated patients and 0.56/100 patient-years in placebo-treated patients after adjusting for prior cholecystectomy. Serious events of acute cholecystitis were reported in 0.5% and 0.3% of patients on TRULICITY and placebo respectively.

Heart Rate Increase and Tachycardia-Related Adverse Reactions

TRULICITY 0.75 mg and 1.5 mg resulted in a mean increase in heart rate (HR) of 2-4 beats per minute (bpm).

Adverse reactions of sinus tachycardia were reported more frequently in patients exposed to TRULICITY. Sinus tachycardia was reported in 3.0%, 2.8%, and 5.6% of patients treated with placebo, TRULICITY 0.75 mg and TRULICITY

1.5 mg, respectively. Persistence of sinus tachycardia (reported at more than 2 visits) was reported in 0.2%, 0.4% and 1.6% of patients treated with placebo, TRULICITY 0.75 mg and TRULICITY 1.5 mg, respectively. Episodes of sinus tachycardia, associated with a concomitant increase from baseline in heart rate of ≥ 15 beats per minute, were reported in 0.7%, 1.3% and 2.2% of patients treated with placebo, TRULICITY 0.75 mg and TRULICITY 1.5 mg, respectively.

Hypersensitivity

Systemic hypersensitivity adverse reactions, sometimes severe (e.g., severe urticaria, systemic rash, facial edema, lip swelling), occurred in 0.5% of patients on TRULICITY in clinical studies.

Injection-site Reactions

In the placebo-controlled studies, injection-site reactions (e.g., injection-site rash, erythema) were reported in 0.5% of TRULICITY-treated patients and in 0.0% of placebo-treated patients.

PR Interval Prolongation and Adverse Reactions of First-Degree Atrioventricular (AV) Block

A mean increase from baseline in PR interval of 2-3 milliseconds was observed in TRULICITY-treated patients in contrast to a mean decrease of 0.9 milliseconds in placebo-treated patients. The adverse reaction of first-degree AV block occurred more frequently in patients treated with TRULICITY than placebo (0.9%, 1.7% and 2.3% for placebo, TRULICITY 0.75 mg and TRULICITY 1.5 mg, respectively). On electrocardiograms, a PR interval increase to at least 220 milliseconds was observed in 0.7%, 2.5% and 3.2% of patients treated with placebo, TRULICITY 0.75 mg and TRULICITY 1.5 mg, respectively.

Amylase and Lipase Increase

Patients exposed to TRULICITY had mean increases from baseline in lipase and/or pancreatic amylase of 14% to 20%, while placebo-treated patients had mean increases of up to 3%.

6.2 Immunogenicity

In clinical studies, 64 (1.6%) TRULICITY-treated patients developed anti-drug antibodies (ADAs) to the active ingredient in TRULICITY (i.e., dulaglutide).

Of the 64 dulaglutide-treated patients that developed dulaglutide ADAs, 34 patients (0.9% of the overall population) had dulaglutide-neutralizing antibodies, and 36 patients (0.9% of the overall population) developed antibodies against native GLP-1.

The detection of antibody formation is highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of antibody (including neutralizing antibody) positivity in an assay may be influenced by several factors including assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, the incidence of antibodies to dulaglutide cannot be directly compared with the incidence of antibodies of other products.

6.3 Postmarketing Experience

The following additional adverse reactions have been reported during post-approval use of TRULICITY. Because these events are reported voluntarily from a population of uncertain size, it is generally not possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- Anaphylactic reactions, angioedema.
- Acute renal failure or worsening of chronic renal failure, sometimes requiring hemodialysis.

7 DRUG INTERACTIONS

7.1 Oral Medications

TRULICITY delays gastric emptying and thus has the potential to reduce the rate of absorption of concomitantly administered oral medications. The delay in gastric emptying is dose-dependent but is attenuated with the recommended dose escalation to higher doses of TRULICITY [see *Dosage and Administration (2.1)*]. The delay is largest after the first dose and diminishes with subsequent doses. In clinical pharmacology studies, TRULICITY 1.5 mg did not affect the absorption of the tested orally administered medications to a clinically relevant degree [see *Clinical Pharmacology (12.3)*]. There is limited experience with the use of concomitant medications in clinical trials with TRULICITY doses of 3 mg and 4.5 mg.

Monitor drug levels of oral medications with a narrow therapeutic index (e.g., warfarin) when concomitantly administered with TRULICITY.

7.2 Concomitant Use with an Insulin Secretagogue (e.g., Sulfonylurea) or with Insulin

When initiating TRULICITY, consider reducing the dose of concomitantly administered insulin secretagogues (such as sulfonylureas) or insulin to reduce the risk of hypoglycemia [see *Warnings and Precautions (5.4)*].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Limited data with TRULICITY in pregnant women are not sufficient to determine a drug-associated risk for major birth defects and miscarriage. There are clinical considerations regarding the risks of poorly controlled diabetes in pregnancy [see *Clinical Considerations*]. Based on animal reproduction studies, there may be risks to the fetus from exposure to dulaglutide during pregnancy. TRULICITY should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

In pregnant rats administered dulaglutide during organogenesis, early embryonic deaths, fetal growth reductions, and fetal abnormalities occurred at systemic exposures at least 6-times human exposure at the maximum recommended human dose (MRHD) of 4.5 mg/week. In pregnant rabbits administered dulaglutide during organogenesis, major fetal abnormalities occurred at 5-times human exposure at the MRHD. Adverse embryo/fetal effects in animals occurred in association with decreased maternal weight and food consumption attributed to the pharmacology of dulaglutide [see *Data*].

The estimated background risk of major birth defects is 6–10% in women with pre-gestational diabetes with an HbA1c >7% and has been reported to be as high as 20–25% in women with an HbA1c >10%. The estimated background risk of miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2–4% and 15–20%, respectively.

Clinical Considerations

Disease-associated maternal and/or embryo/fetal risk

Poorly controlled diabetes in pregnancy increases the maternal risk for diabetic ketoacidosis, pre-eclampsia, spontaneous abortions, preterm delivery and delivery complications. Poorly controlled diabetes increases the fetal risk for major birth defects, stillbirth, and macrosomia-related morbidity.

Data

Animal Data

Pregnant rats given subcutaneous doses of 0.49, 1.63, or 4.89 mg/kg dulaglutide every 3 days during organogenesis had systemic exposures 2-, 6-, and 18-times human exposure at the maximum recommended human dose (MRHD) of 4.5 mg/week, respectively, based on plasma area under the time-concentration curve (AUC) comparison. Reduced fetal weights associated with decreased maternal food intake and decreased weight gain attributed to the pharmacology of dulaglutide were observed at ≥ 1.63 mg/kg. Irregular skeletal ossifications and increases in post-implantation loss also were observed at 4.89 mg/kg.

In pregnant rabbits given subcutaneous doses of 0.04, 0.12, or 0.41 mg/kg dulaglutide every 3 days during organogenesis, systemic exposures in pregnant rabbits were 0.5-, 2-, and 5-times human exposure at the MRHD, based on plasma AUC comparison. Fetal visceral malformation of lung lobular agenesis and skeletal malformations of the vertebrae and/or ribs were observed in conjunction with decreased maternal food intake and decreased weight gain attributed to the pharmacology of dulaglutide at 0.41 mg/kg.

In a prenatal-postnatal study in F₀ maternal rats given subcutaneous doses of 0.2, 0.49, or 1.63 mg/kg every third day from implantation through lactation, systemic exposures in pregnant rats were 1-, 2-, and 7-times human exposure at the MRHD, based on plasma AUC comparison. F₁ pups from F₀ maternal rats given 1.63 mg/kg dulaglutide had statistically significantly lower mean body weight from birth through postnatal day 63 for males and postnatal day 84 for females. F₁ offspring from F₀ maternal rats receiving 1.63 mg/kg dulaglutide had decreased forelimb and hindlimb grip strength and males had delayed balano-preputial separation. Females had decreased startle response. These physical findings may relate to the decreased size of the offspring relative to controls as they appeared at early postnatal assessments but were not observed at a later assessment. F₁ female offspring of the F₀ maternal rats given 1.63 mg/kg of dulaglutide had a longer mean escape time and a higher mean number of errors relative to concurrent control during 1 of 2 trials in the memory evaluation portion of the Biel water maze. These findings occurred in conjunction with decreased F₀ maternal food intake and decreased weight gain attributed to the pharmacologic activity at 1.63 mg/kg. The human relevance of these memory deficits in the F₁ female rats is not known.

8.2 Lactation

Risk Summary

There are no data on the presence of dulaglutide in human milk, the effects on the breastfed infant, or the effects on milk production. The presence of dulaglutide in milk of treated lactating animals was not determined. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for TRULICITY and any potential adverse effects on the breastfed infant from TRULICITY or from the underlying maternal condition.

8.4 Pediatric Use

Safety and effectiveness of TRULICITY have not been established in pediatric patients. TRULICITY is not recommended for use in pediatric patients younger than 18 years.

8.5 Geriatric Use

In the glycemic control trials [see *Clinical Studies (14.1)*], 620 (18.6%) TRULICITY-treated patients were 65 years of age or older and 65 (1.9%) TRULICITY-treated patients were 75 years of age or older at baseline. In the TRULICITY 1.5 mg treatment arm of the REWIND trial [see *Clinical Studies (14.2)*], a total of 2619 (52.9%) patients were 65 years of age or older, and 484 (9.8%) patients were 75 years of age or older at baseline.

No overall differences in safety or efficacy were detected between these patients and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

8.6 Renal Impairment

TRULICITY has been studied in patients with varying degrees of renal function, including a dedicated study in patients with moderate to severe chronic kidney disease. No overall differences in safety or effectiveness were observed in these studies according to renal function [see *Clinical Studies (14)*].

In a clinical pharmacology study in subjects with renal impairment, including end-stage renal disease (ESRD), no clinically relevant change in dulaglutide pharmacokinetics (PK) was observed. In the 52-week study in patients with type 2 diabetes and moderate to severe renal impairment, the PK behavior of TRULICITY 0.75 mg and 1.5 mg once weekly was similar to that demonstrated in previous clinical studies [see *Clinical Pharmacology (12.3)*].

No dose adjustment is recommended in patients with renal impairment including end-stage renal disease (ESRD). Monitor renal function in patients with renal impairment reporting severe adverse gastrointestinal reactions. Use TRULICITY with caution in patients with ESRD [see *Warning and Precautions (5.5)*, *Clinical Pharmacology (12.3)*].

8.7 Hepatic Impairment

In a clinical pharmacology study in subjects with varying degrees of hepatic impairment, no clinically relevant change in dulaglutide PK was observed [see *Clinical Pharmacology (12.3)*]. However, there is limited clinical experience in patients with mild, moderate, or severe hepatic impairment; therefore, use TRULICITY with caution in these patient populations.

8.8 Gastroparesis

Dulaglutide slows gastric emptying. TRULICITY has not been studied in patients with preexisting gastroparesis. Use TRULICITY with caution in patients with gastroparesis.

10 OVERDOSAGE

Overdoses have been reported in clinical studies. Effects associated with these overdoses were primarily mild or moderate gastrointestinal events (e.g., nausea, vomiting) and non-severe hypoglycemia. In the event of overdose, appropriate supportive care (including frequent plasma glucose monitoring) should be initiated according to the patient's clinical signs and symptoms.

11 DESCRIPTION

Dulaglutide is a human glucagon-like peptide-1 (GLP-1) receptor agonist. The molecule is a fusion protein that consists of 2 identical, disulfide-linked chains, each containing an N-terminal GLP-1 analog sequence covalently linked to the Fc portion of a modified human immunoglobulin G4 (IgG4) heavy chain by a small peptide linker and is produced using mammalian cell (Chinese hamster ovary) culture. The GLP-1 analog portion of dulaglutide is 90% homologous to native human GLP-1 (7-37). Structural modifications were introduced in the GLP-1 part of the molecule responsible for interaction with the enzyme dipeptidyl-peptidase-IV (DPP-4). Additional modifications were made in an area with a potential T-cell epitope and in the areas of the IgG4 Fc part of the molecule responsible for binding the high-affinity Fc receptors and half-antibody formation. The overall molecular weight of dulaglutide is approximately 63 kilodaltons.

TRULICITY (dulaglutide) injection is a clear, colorless, sterile, preservative-free solution for subcutaneous use. Each single-dose pen contains a 0.5 mL solution of 0.75 mg, 1.5 mg, 3 mg, or 4.5 mg of dulaglutide and the following excipients: citric acid anhydrous (0.07 mg), mannitol (23.2 mg), polysorbate 80 (0.10 mg for 0.75 mg and 1.5 mg; 0.125 mg for 3 mg and 4.5 mg), and trisodium citrate dihydrate (1.37 mg), in water for injection.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

TRULICITY contains dulaglutide, which is a human GLP-1 receptor agonist with 90% amino acid sequence homology to endogenous human GLP-1 (7-37). Dulaglutide activates the GLP-1 receptor, a membrane-bound cell-surface receptor coupled to adenylyl cyclase in pancreatic beta cells. Dulaglutide increases intracellular cyclic AMP (cAMP) in beta cells leading to glucose-dependent insulin release. Dulaglutide also decreases glucagon secretion and slows gastric emptying.

12.2 Pharmacodynamics

TRULICITY lowers fasting glucose and reduces postprandial glucose (PPG) concentrations in patients with type 2 diabetes mellitus. The reduction in fasting and postprandial glucose can be observed after a single dose.

Fasting and Postprandial Glucose

In a clinical pharmacology study in patients with type 2 diabetes mellitus, treatment with once weekly TRULICITY resulted in a reduction of fasting and 2-hour PPG concentrations, and postprandial serum glucose incremental AUC, when compared to placebo (-25.6 mg/dL, -59.5 mg/dL, and -197 mg*h/dL, respectively); these effects were sustained after 6 weeks of dosing with the 1.5 mg dose.

First- and Second-Phase Insulin Secretion

Both first- and second-phase insulin secretion were increased in patients with type 2 diabetes treated with TRULICITY compared with placebo.

Insulin and Glucagon Secretion

TRULICITY stimulates glucose-dependent insulin secretion and reduces glucagon secretion. Treatment with TRULICITY 0.75 mg and 1.5 mg once weekly increased fasting insulin from baseline at Week 26 by 35.38 and 17.50 pmol/L, respectively, and C-peptide concentration by 0.09 and 0.07 nmol/L, respectively, in a monotherapy study. In the same study, fasting glucagon concentration was reduced by 1.71 and 2.05 pmol/L from baseline with TRULICITY 0.75 mg and 1.5 mg, respectively.

Gastric Motility

Dulaglutide causes a delay of gastric emptying. The delay in gastric emptying is dose-dependent but is attenuated with adequate dose escalation to higher doses of TRULICITY. The delay is largest after the first dose and diminishes with subsequent doses.

Cardiac Electrophysiology (QTc)

The effect of dulaglutide on cardiac repolarization was tested in a thorough QTc study. Dulaglutide did not produce QTc prolongation at doses of 4 and 7 mg. The maximum recommended dose is 4.5 mg once weekly.

12.3 Pharmacokinetics

The pharmacokinetics of dulaglutide is similar between healthy subjects and patients with type 2 diabetes mellitus. Following subcutaneous administration, the time to maximum plasma concentration of dulaglutide at steady state ranges from 24 to 72 hours, with a median of 48 hours. After reaching steady state, the accumulation ratio was approximately 1.56. Steady-state plasma dulaglutide concentrations were achieved between 2 and 4 weeks following once weekly administration. Site of subcutaneous administration (abdomen, upper arm, and thigh) had no statistically significant effect on the exposure to dulaglutide.

Absorption – The mean absolute bioavailability of dulaglutide following subcutaneous administration of single 0.75 mg and 1.5 mg doses was 65% and 47%, respectively. Absolute subcutaneous bioavailability for 3 mg and 4.5 mg doses were estimated to be similar to 1.5 mg although this has not been specifically studied. Dulaglutide concentrations increased approximately proportional to dose from 0.75 mg to 4.5 mg.


Distribution – Apparent population mean central volume of distribution was 3.09 L and the apparent population mean peripheral volume of distribution was 5.98 L.

Metabolism – Dulaglutide is presumed to be degraded into its component amino acids by general protein catabolism pathways.

Elimination –The apparent population mean clearance of dulaglutide was 0.142 L/h. The elimination half-life of dulaglutide was approximately 5 days.

Specific Populations

The intrinsic factors of age, gender, race, ethnicity, body weight, or renal or hepatic impairment do not have a clinically relevant effect on the PK of dulaglutide as shown in Figure 1.



Abbreviations: AUC = area under the time-concentration curve; CI = confidence interval; C_{max} = maximum concentration; ESRD = end-stage renal disease; PK = pharmacokinetics.

Note: Reference values for weight, age, gender, and race comparisons are 93 kg, 56 years old, male, and white, respectively; reference groups for renal and hepatic impairment data are subjects with normal renal and hepatic function from the respective clinical pharmacology studies. The weight values shown in the plot (70 and 120 kg) are the 10th and 90th percentiles of weight in the PK population.

Figure 1: Impact of intrinsic factors on dulaglutide pharmacokinetics.

Renal – Dulaglutide systemic exposure was increased by 20, 28, 14 and 12% for mild, moderate, severe, and ESRD renal impairment sub-groups, respectively, compared to subjects with normal renal function. The corresponding values for increase in C_{max} were 13, 23, 20 and 11%, respectively (Figure 1). Additionally, in a 52 week clinical study in patients with type 2 diabetes and moderate to severe renal impairment, the PK behavior of TRULICITY 0.75 mg and 1.5 mg once weekly was similar to that demonstrated in previous clinical studies [see *Warning and Precautions (5.5)*, *Use in Specific Population (8.6)*].

Hepatic – Dulaglutide systemic exposure decreased by 23, 33 and 21% for mild, moderate and severe hepatic impairment groups, respectively, compared to subjects with normal hepatic function, and C_{max} was decreased by a similar magnitude (Figure 1) [see *Use in Specific Population (8.7)*].


Drug Interactions

The potential effect of co-administered medications on the PK of dulaglutide 1.5 mg and vice versa was studied in several single- and multiple-dose studies in healthy subjects, patients with type 2 diabetes mellitus, and patients with hypertension.

Potential for Dulaglutide to Influence the Pharmacokinetics of Other Drugs

Dulaglutide slows gastric emptying and, as a result, may reduce the extent and rate of absorption of orally co-administered medications. In clinical pharmacology studies, dulaglutide at a dose of 1.5 mg did not affect the absorption of the tested orally administered medications to any clinically relevant degree. The delay in gastric emptying is dose-

dependent but is attenuated with the recommended dose escalation to higher doses of TRULICITY [see *Dosage and Administration (2.1)*, *Drug Interactions (7.1)*]. The delay is largest after the first dose and diminishes with subsequent doses. PK measures indicating the magnitude of these interactions are presented in Figure 2.



Abbreviations: AUC = area under the time-concentration curve; CI = confidence interval; C_{max} = maximum concentration; PK = pharmacokinetics. Note: Reference group is co-administered medication given alone.

Figure 2: Impact of dulaglutide 1.5 mg on the pharmacokinetics of co-administered medications.

Potential for Co-administered Drugs to Influence the Pharmacokinetics of Dulaglutide

In a clinical pharmacology study, the co-administration of a single dose of 1.5 mg dulaglutide with steady-state dose of 100 mg sitagliptin caused an increase in dulaglutide AUC and C_{max} of approximately 38% and 27%, which is not considered clinically relevant.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, and Impairment of Fertility

A 2-year carcinogenicity study was conducted with dulaglutide in male and female rats at doses of 0.05, 0.5, 1.5, and 5 mg/kg (0.2-, 3-, 8-, and 24-fold the MRHD of 4.5 mg once weekly based on AUC) administered by subcutaneous injection twice weekly. In rats, dulaglutide caused a dose-related and treatment-duration-dependent increase in the incidence of thyroid C-cell tumors (adenomas and/or carcinomas) compared to controls, at ≥3-fold the MRHD based on AUC. A statistically significant increase in C-cell adenomas was observed in rats receiving dulaglutide at ≥0.5 mg/kg. Numerical increases in thyroid C-cell carcinomas occurred at 5 mg/kg (24 times the MRHD based on AUC) and were considered to be treatment-related despite the absence of statistical significance.

A 6-month carcinogenicity study was conducted with dulaglutide in rasH2 transgenic mice at doses of 0.3, 1, and 3 mg/kg administered by subcutaneous injection twice weekly. Dulaglutide did not produce increased incidences of thyroid C-cell hyperplasia or neoplasia at any dose.

Dulaglutide is a recombinant protein; no genotoxicity studies have been conducted.

Human relevance of thyroid C-cell tumors in rats is unknown and could not be determined by clinical studies or nonclinical studies [see *Boxed Warning and Warnings and Precautions (5.1)*].

In fertility and early embryonic development studies in male and female rats, no adverse effects of dulaglutide on sperm morphology, mating, fertility, conception, and embryonic survival were observed at up to 16.3 mg/kg (55-fold the MRHD based on AUC). In female rats, an increase in the number of females with prolonged diestrus and a dose-related decrease in the mean number of corpora lutea, implantation sites, and viable embryos were observed at ≥ 4.9 mg/kg (≥ 13 -fold the MRHD based on AUC), which occurred in the presence of decreased maternal food consumption and body weight gain.

13.2 Animal Toxicology and/or Pharmacology

Zucker diabetic fatty (ZDF) rats were given 0.5, 1.5, or 5 mg/kg/twice weekly of dulaglutide (1-, 3-, and 13-fold the MRHD based on AUC) for 3 months. Increases of 12% to 33% in total and pancreatic amylase, but not lipase, were observed at all doses without microscopic pancreatic inflammatory correlates in individual animals. Other changes in the dulaglutide-treated animals included increased interlobular ductal epithelium without active ductal cell proliferation (≥ 0.5 mg/kg), increased acinar atrophy with/without inflammation (≥ 1.5 mg/kg), and increased neutrophilic inflammation of the acinar pancreas (5 mg/kg).

Treatment of monkeys for 12 months with 8.15 mg/kg/twice weekly of dulaglutide (nearly 200-fold the MRHD based on AUC) demonstrated no evidence of pancreatic inflammation or pancreatic intraepithelial neoplasia. In 4 of 19 monkeys on dulaglutide treatment, there was an increase in goblet cells within the pancreatic ducts, but no differences from the control group in total amylase or lipase at study termination. There were no proliferative changes in the thyroid C-cells.

14 CLINICAL STUDIES

14.1 Glycemic Control Trials in Adults with Type 2 Diabetes Mellitus

TRULICITY has been studied as monotherapy and in combination with metformin, sulfonylurea, metformin and sulfonylurea, metformin and thiazolidinedione, sodium-glucose co-transporter-2 inhibitors (SGLT2i) with or without metformin, basal insulin with or without metformin, and prandial insulin with or without metformin. TRULICITY has also been studied in patients with type 2 diabetes mellitus and moderate to severe renal impairment.

Dose escalation was performed in one study with TRULICITY doses up to 4.5 mg added to metformin. All other clinical studies evaluated TRULICITY 0.75 mg and 1.5 mg without dose escalation; patients were initiated and maintained on either 0.75 mg or 1.5 mg for the duration of the trials.

In patients with type 2 diabetes mellitus, TRULICITY produced reductions from baseline in HbA1c compared to placebo. No overall differences in glycemic effectiveness were observed across demographic subgroups (age, gender, race/ethnicity, duration of diabetes).

Monotherapy

In a double-blind study with primary endpoint at 26 weeks, 807 patients inadequately treated with diet and exercise, or with diet and exercise and one antidiabetic agent used at submaximal dose, were randomized to TRULICITY 0.75 mg once weekly, TRULICITY 1.5 mg once weekly, or metformin 1500 to 2000 mg/day following a two-week washout. Seventy-five percent (75%) of the randomized population were treated with one antidiabetic agent at the screening visit. Most patients previously treated with an antidiabetic agent were receiving metformin (~90%) at a median dose of 1000 mg daily and approximately 10% were receiving a sulfonylurea.

Patients had a mean age of 56 years and a mean duration of type 2 diabetes of 3 years. Forty-four percent were male. The White, Black and Asian race accounted for 74%, 7% and 8% of the population, respectively. Twenty-nine percent of the study population were from the US.

Treatment with TRULICITY 0.75 mg and 1.5 mg once weekly resulted in reduction in HbA1c from baseline at 26-weeks (Table 4). The difference in observed effect size between TRULICITY 0.75 mg and 1.5 mg, respectively, and metformin excluded the pre-specified non-inferiority margin of 0.4%.

Table 4: Results at Week 26 in a Trial of TRULICITY as Monotherapy^a

	26-Week Primary Time Point		
	TRULICITY 0.75 mg	TRULICITY 1.5 mg	Metformin 1500-2000 mg
Intent-to-Treat (ITT) Population (N)[‡]	270	269	268
HbA1c (%) (Mean)			
Baseline	7.6	7.6	7.6
Change from baseline ^b	-0.7	-0.8	-0.6

Fasting Serum Glucose (mg/dL) (Mean)			
Baseline	161	164	161
Change from baseline ^b	-26	-29	-24
Body Weight (kg) (Mean)			
Baseline	91.8	92.7	92.4
Change from baseline ^b	-1.4	-2.3	-2.2

Abbreviation: HbA1c = hemoglobin A1c.

^a Intent-to-treat population. Last observation carried forward (LOCF) was used to impute missing data. Data post-onset of rescue therapy are treated as missing. At Week 26, primary efficacy was missing for 10%, 12% and 14% of individuals randomized to TRULICITY 0.75 mg, TRULICITY 1.5 mg and metformin, respectively.

^b Least-squares mean adjusted for baseline value and other stratification factors.

[†] Subjects included in the analysis are a subset of the ITT population that had at least one post-baseline assessment. The primary analysis included 265 individuals in each of the treatment arms.

Combination Therapy

Add-on to Metformin

In this placebo-controlled, double-blind study with primary endpoint at 52 weeks, 972 patients were randomized to placebo, TRULICITY 0.75 mg once weekly, TRULICITY 1.5 mg once weekly, or sitagliptin 100 mg/day (after 26 weeks, patients in the placebo treatment group received blinded sitagliptin 100 mg/day for the remainder of the study), all as add-on to metformin. Randomization occurred after an 11-week lead-in period to allow for a metformin titration period, followed by a 6-week glycemic stabilization period. Patients had a mean age of 54 years; mean duration of type 2 diabetes of 7 years; 48% were male; race: White, Black and Asian were 53%, 4% and 27%, respectively; and 24% of the study population were in the US.

At the 26-week placebo-controlled time point, the HbA1c change was 0.1%, -1.0%, -1.2%, and -0.6% for placebo, TRULICITY 0.75 mg, TRULICITY 1.5 mg, and sitagliptin, respectively. The percentage of patients who achieved HbA1c <7.0% was 22%, 56%, 62% and 39% for placebo, TRULICITY 0.75 mg, TRULICITY 1.5 mg, and sitagliptin, respectively. At 26 weeks, there was a mean weight reduction of 1.4 kg, 2.7 kg, 3.0 kg, and 1.4 kg for placebo, TRULICITY 0.75 mg, TRULICITY 1.5 mg, and sitagliptin, respectively. There was a mean reduction of fasting glucose of 9 mg/dL, 35 mg/dL, 41 mg/dL, and 18 mg/dL for placebo, TRULICITY 0.75 mg, TRULICITY 1.5 mg, and sitagliptin, respectively.

Treatment with TRULICITY 0.75 mg and 1.5 mg once weekly resulted in a statistically significant reduction in HbA1c compared to placebo (at 26 weeks) and compared to sitagliptin (at 26 and 52 weeks), all in combination with metformin (Table 5 and Figure 3).

Table 5: Results at Week 52 of TRULICITY Compared to Sitagliptin used as Add-On to Metformin^a

	52-Week Primary Time Point		
	TRULICITY 0.75 mg	TRULICITY 1.5 mg	Sitagliptin 100 mg
Intent-to-Treat (ITT) Population (N)[†]	281	279	273
HbA1c (%) (Mean)			
Baseline	8.2	8.1	8.0
Change from baseline ^b	-0.9	-1.1	-0.4
Difference from sitagliptin ^b (95% CI)	-0.5 (-0.7, -0.3) ^{††}	-0.7 (-0.9, -0.5) ^{††}	-
Percentage of patients HbA1c <7.0%	49 ^{###}	59 ^{###}	33
Fasting Plasma Glucose (mg/dL) (Mean)			
Baseline	174	173	171
Change from baseline ^b	-30	-41	-14
Difference from sitagliptin ^b (95% CI)	-15 (-22, -9)	-27 (-33, -20)	-
Body Weight (kg) (Mean)			
Baseline	85.5	86.5	85.8
Change from baseline ^b	-2.7	-3.1	-1.5

Difference from sitagliptin ^b (95% CI)	-1.2 (-1.8, -0.6)	-1.5 (-2.1, -0.9)	-
---	-------------------	-------------------	---

Abbreviations: HbA1c = hemoglobin A1c.


^a All ITT patients randomized after the dose-finding portion of the study. Last observation carried forward (LOCF) was used to impute missing data. At Week 52 primary efficacy was missing for 15%, 19%, and 20% of individuals randomized to TRULICITY 0.75 mg, TRULICITY 1.5 mg and sitagliptin, respectively.

^b Least-squares (LS) mean adjusted for baseline value and other stratification factors.

[†] Subjects included in the analysis are a subset of the ITT population that had at least one post-baseline assessment. The primary analysis included 276, 277, and 270 individuals randomized to TRULICITY 0.75 mg, TRULICITY 1.5 mg and sitagliptin, respectively.

^{††} Multiplicity adjusted 1-sided p-value <0.001, for superiority of TRULICITY compared to sitagliptin, assessed only for HbA1c.

^{##} p<0.001 TRULICITY compared to sitagliptin, assessed only for HbA1c <7.0%.



Number of subjects with observed data

Placebo	139	108	
TRULICITY 0.75 mg	281	258	238
TRULICITY 1.5 mg	279	249	225
Sitagliptin	273	241	219

Mean HbA1c adjusted for baseline HbA1c and country.

Figure 3: Adjusted Mean HbA1c at each Time Point (ITT, MMRM) and at Week 52 (ITT, LOCF)

TRULICITY 3 mg and 4.5 mg Add-on to Metformin

In this parallel-arm, double-blind study with primary endpoint at 36 weeks, a total of 1842 patients were randomized 1:1:1 to TRULICITY 1.5 mg, TRULICITY 3 mg, or TRULICITY 4.5 mg once weekly, all as add-on to metformin (NCT03495102).

Following randomization, all patients received TRULICITY 0.75 mg once weekly. The dose was increased every 4 weeks to the next higher dose until the patients reached the assigned study dose (1.5 mg, 3 mg, or 4.5 mg). Patients were to remain on the assigned study dose for the duration of the study.

Patients had a mean age of 57.1 years; a mean duration of type 2 diabetes of 7.6 years; 51.2% were male; race: White, Black, and Asian were 85.8%, 4.5%, and 2.4%, respectively; and 27.6% of the study population was in the US.

At 36 weeks, treatment with TRULICITY 4.5 mg resulted in a statistically significant reduction in HbA1c and in body weight compared to TRULICITY 1.5 mg (Table 6 and Figure 4).

Table 6. Results at Week 36 of TRULICITY 1.5 mg Compared to 3 mg and 4.5 mg as Add-On to Metformin^a

	36-Week Primary Time Point		
	TRULICITY 1.5 mg	TRULICITY 3 mg	TRULICITY 4.5 mg
Intent-to-Treat (ITT) Population (N)	612	616	614
HbA1c (%) (Mean)			
Baseline	8.6	8.6	8.6
Change from baseline ^b	-1.5	-1.6	-1.8
Difference from 1.5 mg ^b (95% CI)		-0.1 (-0.2, 0.0)	-0.2 (-0.4, -0.1) [^]
Percentage of patients HbA1c <7.0%^c	50	56	62
Fasting Serum Glucose (mg/dL) (Mean)			
Baseline	185	184	183
Change from baseline ^b	-45	-46	-51
Difference from 1.5 mg ^b (95% CI)		- 2 (-7, 3)	-6 (-11, -2)
Body Weight (kg) (Mean)			
Baseline	95.5	96.3	95.4
Change from baseline ^b	-3.0	-3.8	-4.6
Difference from 1.5 mg ^b (95% CI)		-0.9 (-1.4, -0.4)	-1.6 (-2.2, -1.1) ^{^^}

Abbreviations: HbA1c = hemoglobin A1c


^a Intent-to-treat population. At Week 36, primary efficacy was missing for 7%, 7%, and 6% of individuals treated with TRULICITY 1.5 mg, TRULICITY 3 mg, and TRULICITY 4.5 mg, respectively.

^b Least-squares mean adjusted for baseline value and other stratification factors. Missing data were imputed using multiple imputation.

^c Patients with missing HbA1c data at Week 36 were considered as not achieving HbA1c target.

[^] p=0.0001 for superiority compared to TRULICITY 1.5 mg, overall type I error controlled.

^{^^} p<0.0001 for superiority compared to TRULICITY 1.5 mg, overall type I error controlled.



Number of subjects with observed data

TRULICITY 1.5 mg	612	567
TRULICITY 3 mg	616	572
TRULICITY 4.5 mg	614	575

Observed mean HbA1c at scheduled visits and retrieved dropout multiple imputation (MI) based estimate at week 36.

Figure 4: Mean HbA1c at each Time Point (ITT) and at Week 36 (ITT, MI)

Add-on to Sulfonylurea

In this 24-week placebo-controlled, double-blind study, 299 patients were randomized to and received placebo or once weekly TRULICITY 1.5 mg, both as add-on to glimepiride. Patients had a mean age of 58 years; mean duration of type 2 diabetes of 8 years; 44% were male; race: White, Black, and Asian were 83%, 4%, and 2%, respectively; and 24% of the study population were in the US.

At 24 weeks, treatment with once weekly TRULICITY 1.5 mg resulted in a statistically significant reduction in HbA1c compared to placebo (Table 7).

Table 7: Results at Week 24 of TRULICITY Compared to Placebo as Add-On to Glimepiride^a

	24-Week Primary Time Point	
	Placebo	TRULICITY 1.5 mg
Intent-to-Treat (ITT) Population (N)	60	239
HbA1c (%) (Mean)		
Baseline	8.4	8.4
Change from baseline ^b	-0.3	-1.3
Difference from placebo ^b (95% CI)		-1.1 (-1.4, -0.7) ^{††}
Percentage of patients HbA1c <7.0%^c	17	50 ^{††}
Fasting Serum Glucose (mg/dL) (Mean)		
Baseline	175	178
Change from baseline ^b	2	-28
Difference from placebo ^b (95% CI)		-30 (-44, -15) ^{††}

Body Weight (kg) (Mean)		
Baseline	89.5	84.5
Change from baseline ^b	-0.2	-0.5
Difference from placebo ^b (95% CI)		-0.4 (-1.2, 0.5)

Abbreviations: HbA1c = hemoglobin A1c.

^a Intent-to-treat population. Data post-onset of rescue therapy are treated as missing. At Week 24 primary efficacy was missing for 10% and 12% of individuals randomized to TRULICITY 1.5 mg and placebo, respectively.

^b Least-squares mean from ANCOVA adjusted for baseline value and other stratification factors. Placebo multiple imputation, with respect to the baseline values, was used to model a wash-out of the treatment effect for subjects having missing Week 24 data.

^c Patients with missing HbA1c data at Week 24 were considered as non-responders.

^{††} $p < 0.001$ for superiority of TRULICITY 1.5 mg compared to placebo, overall type I error controlled.

Add-on to Metformin and Thiazolidinedione

In this placebo-controlled study with primary endpoint at 26 weeks, 976 patients were randomized to and received placebo, TRULICITY 0.75 mg once weekly, TRULICITY 1.5 mg once weekly, or exenatide 10 mcg BID, all as add-on to maximally tolerated doses of metformin (≥ 1500 mg per day) and pioglitazone (up to 45 mg per day). Exenatide treatment group assignment was open-label while the treatment assignments to placebo, TRULICITY 0.75 mg, and TRULICITY 1.5 mg were blinded. After 26 weeks, patients in the placebo treatment group were randomized to either TRULICITY 0.75 mg once weekly or TRULICITY 1.5 mg once weekly to maintain study blind. Randomization occurred after a 12-week lead-in period; during the initial 4 weeks of the lead-in period, patients were titrated to maximally tolerated doses of metformin and pioglitazone; this was followed by an 8-week glycemic stabilization period prior to randomization. Patients randomized to exenatide started at a dose of 5 mcg BID for 4 weeks and then were escalated to 10 mcg BID. Patients had a mean age of 56 years; mean duration of type 2 diabetes of 9 years; 58% were male; race: White, Black and Asian were 74%, 8% and 3%, respectively; and 81% of the study population were in the US.

Treatment with TRULICITY 0.75 mg and 1.5 mg once weekly resulted in a statistically significant reduction in HbA1c compared to placebo (at 26 weeks) and compared to exenatide at 26 weeks (Table 8 and Figure 5). Over the 52-week study period, the percentage of patients who required glycemic rescue was 8.9% in the TRULICITY 0.75 mg once weekly + metformin and pioglitazone treatment group, 3.2% in the TRULICITY 1.5 mg once weekly + metformin and pioglitazone treatment group, and 8.7% in the exenatide BID + metformin and pioglitazone treatment group.

Table 8: Results at Week 26 of TRULICITY Compared to Placebo and Exenatide, All as Add-On to Metformin and Thiazolidinedione^a

	26-Week Primary Time Point			
	Placebo	TRULICITY 0.75 mg	TRULICITY 1.5 mg	Exenatide 10 mcg BID
Intent-to-Treat (ITT) Population (N)[†]	141	280	279	276
HbA1c (%) (Mean)				
Baseline	8.1	8.1	8.1	8.1
Change from baseline ^b	-0.5	-1.3	-1.5	-1.0
Difference from placebo ^b (95% CI)	-	-0.8 (-1.0, -0.7) ^{††}	-1.1 (-1.2, -0.9) ^{††}	-
Difference from exenatide ^b (95% CI)	-	-0.3 (-0.4, -0.2) ^{††}	-0.5 (-0.7, -0.4) ^{††}	-
Percentage of patients HbA1c <7.0%	43	66 ^{**} , ^{##}	78 ^{**} , ^{##}	52
Fasting Serum Glucose (mg/dL) (Mean)				
Baseline	166	159	162	164
Change from baseline ^b	-5	-34	-42	-24
Difference from placebo ^b (95% CI)	-	-30 (-36, -23)	-38 (-45, -31)	-
Difference from exenatide ^b (95% CI)	-	-10 (-15, -5)	-18 (-24, -13)	-

Body Weight (kg) (Mean)

Baseline	94.1	95.5	96.2	97.4
Change from baseline ^b	1.2	0.2	-1.3	-1.1
Difference from placebo ^b (95% CI)	-	-1.0 (-1.8, -0.3)	-2.5 (-3.3, -1.8)	-
Difference from exenatide ^b (95% CI)	-	1.3 (0.6, 1.9)	-0.2 (-0.9, 0.4)	-

Abbreviations: BID = twice daily; HbA1c = hemoglobin A1c.

^a Intent-to-treat population. Last observation carried forward (LOCF) was used to impute missing data. Data post-onset of rescue therapy are treated as missing. At Week 26, primary efficacy was missing for 23%, 10%, 7% and 12% of individuals randomized to placebo, TRULICITY 0.75 mg, TRULICITY 1.5 mg, and exenatide, respectively.

^b Least-squares (LS) mean adjusted for baseline value and other stratification factors.


[†] Subjects included in the analysis are a subset of the ITT population that had at least one post-baseline assessment. The primary analysis included 119, 269, 271 and 266 individuals randomized to placebo, TRULICITY 0.75 mg, TRULICITY 1.5 mg, and exenatide, respectively.

^{††} Multiplicity adjusted 1-sided p-value <0.001, for superiority of TRULICITY compared to placebo, assessed only for HbA1c.

^{†††} Multiplicity adjusted 1-sided p-value <0.001, for superiority of TRULICITY compared to exenatide, assessed only for HbA1c.

^{**} p<0.001 TRULICITY compared to placebo, assessed only for HbA1c <7.0%.

^{###} p<0.001 TRULICITY compared to exenatide, assessed only for HbA1c <7.0%.



Number of subjects with observed data

Placebo	141	108
TRULICITY 0.75 mg	280	251
TRULICITY 1.5 mg	279	259
Exenatide	276	242

Mean HbA1c adjusted for baseline HbA1c and country.

Figure 5: Adjusted Mean HbA1c at Each Time Point (ITT, MMRM) and at Week 26 (ITT, LOCF)

Combination Therapy with SGLT2i, with or without Metformin

In this 24-week placebo-controlled, double-blind study, 423 patients were randomized to and received TRULICITY 0.75 mg, TRULICITY 1.5 mg, or placebo, as add-on to sodium-glucose co-transporter 2 inhibitor (SGLT2i) therapy (96% with and 4% without metformin). Trulicity was administered once weekly, and SGLT2i was administered according to the local country label. Patients had a mean age of 57 years; mean duration of type 2 diabetes of 9.4 years; 50% were male; race: White, Black, and Asian were 89%, 3%, and 0.2%, respectively; and 21% of the study population was in the US.

At 24 weeks, treatment with once weekly TRULICITY 0.75 mg and 1.5 mg resulted in a statistically significant reduction from baseline in HbA1c compared to placebo (Table 9).

The mean baseline body weight was 90.5, 91.1, and 92.9 kg in the placebo, TRULICITY 0.75 mg, and TRULICITY 1.5 mg groups, respectively. The mean changes from baseline in body weight at Week 24 were -2.0, -2.5, and -2.9 kg for placebo, TRULICITY 0.75 mg, and TRULICITY 1.5 mg, respectively. The difference from placebo (95% CI) was -0.9 kg (-1.7, -0.1) for TRULICITY 1.5 mg.

Table 9: Results at Week 24 of TRULICITY as Add-on to SGLT2i^a

	24-Week Primary Time Point		
	Placebo	TRULICITY 0.75 mg	TRULICITY 1.5 mg
Intent-to-Treat (ITT) Population (N)	140	141	142
HbA1c (%) (Mean)			
Baseline	8.1	8.1	8.0
Change from baseline ^b	-0.6	-1.2	-1.3
Difference from placebo ^b (95% CI)	-	-0.7 (-0.8, -0.5) ^{††}	-0.8 (-0.9, -0.6) ^{††}
Percentage of patients HbA1c <7.0%^c	31	59 ^{††}	67 ^{††}
Fasting Serum Glucose (mg/dL) (Mean)			
Baseline	153	162	161
Change from baseline ^b	-6	-25	-30
Difference from placebo ^b (95% CI)	-	-19 (-25, -13)	-24 (-30, -18) ^{††}

Abbreviations: HbA1c = hemoglobin A1c; SGLT2i = sodium-glucose co-transporter-2 inhibitors.

^a Intent-to-treat population. At Week 24, primary efficacy was missing for 3%, 4%, and 6% of individuals treated with placebo, TRULICITY 0.75 mg, and TRULICITY 1.5 mg, respectively.

^b Least-squares mean adjusted for baseline value and other stratification factors. Placebo multiple imputation, using baseline and 24-week values from the placebo arm, was applied to model a washout of the treatment effect for patients missing 24-week values (HbA1c, fasting serum glucose, and body weight).

^c Patients with missing HbA1c data at Week 24 were considered as non-responders.

^{††} p<0.001 for superiority of TRULICITY compared to placebo, overall type I error controlled.

Add-on to Metformin and Sulfonylurea

In this open-label comparator study (double-blind with respect to TRULICITY dose assignment) with primary endpoint at 52 weeks, 807 patients were randomized to and received TRULICITY 0.75 mg once weekly, TRULICITY 1.5 mg once weekly, or insulin glargine once daily, all as add-on to maximally tolerated doses of metformin and glimepiride. Randomization occurred after a 10-week lead-in period; during the initial 2 weeks of the lead-in period, patients were titrated to maximally tolerated doses of metformin and glimepiride. This was followed by a 6- to 8-week glycemic stabilization period prior to randomization.

Patients randomized to insulin glargine were started on a dose of 10 units once daily. Insulin glargine dose adjustments occurred twice weekly for the first 4 weeks of treatment based on self-measured fasting plasma glucose (FPG), followed by once weekly titration through Week 8 of study treatment, utilizing an algorithm that targeted a fasting plasma glucose of <100 mg/dL. Only 24% of patients were titrated to goal at the 52-week primary endpoint. The dose of glimepiride could be reduced or discontinued after randomization (at the discretion of the investigator) in the event of persistent hypoglycemia. The dose of glimepiride was reduced or discontinued in 28%, 32%, and 29% of patients randomized to TRULICITY 0.75 mg, TRULICITY 1.5 mg, and glargine.

Patients had a mean age of 57 years; mean duration of type 2 diabetes of 9 years; 51% were male; race: White, Black and Asian were 71%, 1% and 17%, respectively; and 0% of the study population were in the US.

Treatment with TRULICITY once weekly resulted in a reduction in HbA1c from baseline at 52 weeks when used in combination with metformin and sulfonylurea (Table 10). The difference in observed effect size between TRULICITY 0.75 mg and 1.5 mg, respectively, and glargine in this trial excluded the pre-specified non-inferiority margin of 0.4%.

Table 10: Results at Week 52 of TRULICITY Compared to Insulin Glargine, Both as Add-on to Metformin and Sulfonylurea^a

	52-Week Primary Time Point		
	TRULICITY 0.75 mg	TRULICITY 1.5 mg	Insulin Glargine
Intent-to-Treat (ITT) Population (N)[‡]	272	273	262
HbA1c (%) (Mean)			
Baseline	8.1	8.2	8.1
Change from baseline ^b	-0.8	-1.1	-0.6
Fasting Serum Glucose (mg/dL) (Mean)			
Baseline	161	165	163
Change from baseline ^b	-16	-27	-32
Difference from insulin glargine ^b (95% CI)	16 (9, 23)	5 (-2, 12)	-
Body Weight (kg) (Mean)			
Baseline	86.4	85.2	87.6
Change from baseline ^b	-1.3	-1.9	1.4
Difference from insulin ^b (95% CI)	-2.8 (-3.4, -2.2)	-3.3 (-3.9, -2.7)	-

Abbreviations: HbA1c = hemoglobin A1c.

^a Intent-to-treat population. Last observation carried forward (LOCF) was used to impute missing data. Data post-onset of rescue therapy are treated as missing. At Week 52, primary efficacy was missing for 17%, 13% and 12% of individuals randomized to TRULICITY 0.75 mg, TRULICITY 1.5 mg and glargine, respectively.

^b Least-squares (LS) mean adjusted for baseline value and other stratification factors.

[‡] Subjects included in the analysis are a subset of the ITT population that had at least one post-baseline assessment. The primary analysis included 267, 263 and 259 individuals randomized to TRULICITY 0.75 mg, TRULICITY 1.5 mg, and glargine, respectively.

Combination Therapy with Basal Insulin, with or without Metformin

In this 28-week placebo-controlled, double-blind study, 300 patients were randomized to placebo or once weekly TRULICITY 1.5 mg, as add-on to titrated basal insulin glargine (with or without metformin). Patients had a mean age of 60 years; mean duration of type 2 diabetes of 13 years; 58% were male; race: White, Black, and Asian were 94%, 4%, and 0.3%, respectively; and 20% of the study population was in the US.

The mean starting dose of insulin glargine was 37 units/day for patients receiving placebo and 41 units/day for patients receiving TRULICITY 1.5 mg. At randomization, the initial insulin glargine dose in patients with HbA1c <8.0% was reduced by 20%.

At 28 weeks, treatment with once weekly TRULICITY 1.5 mg resulted in a statistically significant reduction in HbA1c compared to placebo (Table 11).

Table 11: Results at Week 28 of TRULICITY Compared to Placebo as Add-On to Basal Insulin^a

	28-Week Primary Time Point	
	Placebo	TRULICITY 1.5 mg
Intent-to-Treat (ITT) Population (N)	150	150
HbA1c (%) (Mean)		
Baseline	8.3	8.4
Change from baseline ^b	-0.7	-1.4
Difference from placebo ^b (95% CI)		-0.7 (-0.9, -0.5) ^{††}
Percentage of patients HbA1c <7.0%^c	33	67 ^{††}

Fasting Serum Glucose (mg/dL) (Mean)		
Baseline	156	157
Change from baseline ^b	-30	-44
Difference from placebo ^b (95% CI)		-14 (-23, -4) [†]
Body Weight (kg) (Mean)		
Baseline	92.6	93.3
Change from baseline ^b	0.8	-1.3
Difference from placebo ^b (95% CI)		-2.1 (-2.9, -1.4) ^{††}

Abbreviations: HbA1c = hemoglobin A1c.

^a Intent-to-treat population. At Week 28, primary efficacy was missing for 12% and 8% of individuals randomized to placebo and TRULICITY 1.5 mg, respectively.

^b Least-squares mean from ANCOVA adjusted for baseline value and other stratification factors. Placebo multiple imputation, with respect to baseline values, was used to model a wash-out of the treatment effect for subjects having missing Week 28 data.

^c Patients with missing HbA1c data at Week 28 were considered as non-responders.

^{††} p<0.001 for superiority of TRULICITY 1.5 mg compared to placebo, overall type I error controlled.

[†] p≤0.005 for superiority of TRULICITY 1.5 mg compared to placebo, overall type I error controlled.

Combination Therapy with Prandial Insulin, with or without Metformin

In this open-label comparator study (double-blind with respect to TRULICITY dose assignment) with primary endpoint at 26 weeks, 884 patients on 1 or 2 insulin injections per day were enrolled. Randomization occurred after a 9-week lead-in period; during the initial 2 weeks of the lead-in period, patients continued their pre-study insulin regimen but could be initiated and/or up-titrated on metformin, based on investigator discretion; this was followed by a 7-week glycaemic stabilization period prior to randomization.

At randomization, patients discontinued their pre-study insulin regimen and were randomized to TRULICITY 0.75 mg once weekly, TRULICITY 1.5 mg once weekly, or insulin glargine once daily, all in combination with prandial insulin lispro 3 times daily, with or without metformin. Insulin lispro was titrated in each arm based on preprandial and bedtime glucose, and insulin glargine was titrated to a fasting plasma glucose goal of <100 mg/dL. Only 36% of patients randomized to glargine were titrated to the fasting glucose goal at the 26-week primary timepoint.

Patients had a mean age of 59 years; mean duration of type 2 diabetes of 13 years; 54% were male; race: White, Black and Asian were 79%, 10% and 4%, respectively; and 33% of the study population were in the US.

Treatment with TRULICITY 0.75 mg and 1.5 mg once weekly resulted in a reduction in HbA1c from baseline. The difference in observed effect size between TRULICITY 0.75 mg and 1.5 mg, respectively, and glargine in this trial excluded the pre-specified non-inferiority margin of 0.4% (Table 12).

Table 12: Results at Week 26 of TRULICITY Compared to Insulin Glargine, Both in Combination with Insulin Lispro^a

	26-Week Primary Time Point		
	TRULICITY 0.75 mg	TRULICITY 1.5 mg	Insulin Glargine
Intent-to-Treat (ITT) Population (N)[‡]	293	295	296
HbA1c (%) (Mean)			
Baseline	8.4	8.5	8.5
Change from baseline ^b	-1.6	-1.6	-1.4
Fasting Serum Glucose (mg/dL) (Mean)			
Baseline	150	157	154
Change from baseline ^b	4	-5	-28
Difference from insulin glargine ^b (95% CI)	32 (24, 41)	24 (15, 32)	-
Body Weight (kg) (Mean)			
Baseline	91.7	91.0	90.8
Change from baseline ^b	0.2	-0.9	2.3
Difference from insulin glargine ^b (95% CI)	-2.2 (-2.8, -1.5)	-3.2 (-3.8, -2.6)	-

Abbreviation: HbA1c = hemoglobin A1c

^a Intent-to-treat population. Last observation carried forward (LOCF) was used to impute missing data. Data post-onset of rescue therapy are treated as missing. At Week 26, primary efficacy was missing for 14%, 15%, and 14% of individuals randomized to TRULICITY 0.75 mg, TRULICITY 1.5 mg and glargine, respectively.

^b Least-squares (LS) mean adjusted for baseline value and other stratification factors.

[†] Subjects included in the analysis are a subset of the ITT population that had at least one post-baseline assessment. The primary analysis included 275, 273 and 276 individuals randomized to TRULICITY 0.75 mg, TRULICITY 1.5 mg, and glargine, respectively.

Moderate to Severe Chronic Kidney Disease

In this open-label comparator study (double-blind with respect to TRULICITY dose assignment) with primary endpoint at 26 weeks, a total of 576 patients with type 2 diabetes were randomized and treated to compare TRULICITY 0.75 mg and 1.5 mg with insulin glargine (NCT01621178).

Patients on insulin and other antidiabetic therapy (e.g., oral antidiabetic drugs, pramlintide) had non-insulin therapies discontinued and had their insulin dose adjusted for 12 weeks prior to randomization. Patients on insulin therapy alone maintained a stable insulin dose for 3 weeks prior to randomization. At randomization, patients discontinued their pre-study insulin regimen and patients were randomized to TRULICITY 0.75 mg once weekly, TRULICITY 1.5 mg once weekly, or insulin glargine once daily, all in combination with prandial insulin lispro. For patients randomized to insulin glargine, the initial insulin glargine dose was based on the basal insulin dose prior to randomization. Insulin glargine was allowed to be titrated with a fasting plasma glucose goal of ≤ 150 mg/dL. Insulin lispro was allowed to be titrated with a preprandial and bedtime glucose goal of ≤ 180 mg/dL.

Patients had a mean age of 65 years; a mean duration of type 2 diabetes of 18 years; 52% were male; race: White, Black, and Asian were 69%, 16%, and 3%, respectively; and 32% of the study population were in the US. At baseline, overall mean eGFR was 38 mL/min/1.73 m², 30% of patients had eGFR < 30 mL/min/1.73 m², and 45% of patients had macroalbuminuria. Patients on over 70 units/day of basal insulin were excluded from the study.

Treatment with TRULICITY 0.75 mg and 1.5 mg once weekly resulted in a reduction in HbA1c at 26-weeks from baseline. The difference in observed effect size between TRULICITY 0.75 mg and 1.5 mg, respectively, and glargine in this trial excluded the pre-specified non-inferiority margin of 0.4%. Mean fasting plasma glucose increased in the TRULICITY arms (Table 13).

Mean baseline body weight was 90.9 kg, 88.1 kg, and 88.2 kg in the TRULICITY 0.75 mg, TRULICITY 1.5 mg, and insulin glargine arms, respectively. The mean changes from baseline at Week 26 were -1.1, -2, and 1.9 kg in the TRULICITY 0.75 mg, TRULICITY 1.5 mg, and insulin glargine arms, respectively.

Table 13: Results at Week 26 of TRULICITY Compared to Insulin Glargine, Both in Combination with Insulin Lispro, in Patients with Moderate to Severe Chronic Kidney Disease^a

	26-Week Primary Time Point		
	TRULICITY 0.75 mg	TRULICITY 1.5 mg	Insulin Glargine
Intent-to-Treat Population (N)	190	192	194
HbA1c (%) (Mean)			
Baseline	8.6	8.6	8.6
Change from baseline ^b	-0.9	-1.0	-1.0
Difference from insulin glargine ^b (95% CI)	0.0 (-0.2, 0.3)	-0.1 (-0.3, 0.2)	
Percentage of patients HbA1c <8.0%	73	75	74
Fasting Serum Glucose (mg/dL) (Mean)			
Baseline	167	161	170
Change from baseline ^b	6	14	-23
Difference from insulin glargine ^b (95% CI)	30 (16, 43)	37 (24, 50)	

Abbreviation: HbA1c = hemoglobin A1c

^a Intent-to-treat population (all randomized and treated subjects) was used in the analysis regardless of discontinuation of study drug or initiation of rescue therapy. At Week 26, primary efficacy was missing for 12%, 15%, and 9% of individuals randomized to and treated with TRULICITY 0.75 mg, TRULICITY 1.5 mg, and insulin glargine, respectively. Missing data were imputed using multiple imputation within treatment group.

^b Least-squares (LS) mean from ANCOVA pattern mixture model adjusted for baseline value and other stratification factors.

14.2 Cardiovascular Outcomes Trial in Adults with Type 2 Diabetes Mellitus and Cardiovascular Disease or Multiple Cardiovascular Risk Factors

The REWIND trial (NCT01394952) was a multi-national, multi-center, randomized, placebo-controlled, double-blind trial. In this study, 9901 adult patients with type 2 diabetes mellitus and established cardiovascular (CV) disease or multiple cardiovascular risk factors were randomized to TRULICITY 1.5 mg or placebo both added to standard of care. The median follow-up duration was 5.4 years. The primary endpoint was the time to the first occurrence of a composite 3-component Major Adverse Cardiovascular Events (MACE) outcome, which included CV death, non-fatal myocardial infarction (MI), and non-fatal stroke.

Patients eligible to enter the trial were 50 years of age or older who had type 2 diabetes mellitus, had an HbA1c value $\leq 9.5\%$ with no lower limit at screening, and had either established CV disease, or did not have established CV disease but had multiple CV risk factors. Patients who were confirmed to have established CV disease (31.5% of randomized patients) had a history of at least one of the following: MI (16.2%); myocardial ischemia by a stress test or with cardiac imaging (9.3%); ischemic stroke (5.3%); coronary, carotid, or peripheral artery revascularization (18.0%); unstable angina (5.9%); or hospitalization for unstable angina with at least one of the following: ECG changes, myocardial ischemia on imaging, or a need for percutaneous coronary intervention (12.0%). Patients confirmed to be without established CV disease, but with multiple CV risk factors, comprised 62.8% of the randomized trial population.

At baseline, demographic and disease characteristics were balanced between treatment groups. Patients had a mean age of 66 years; 46% were female; race: White, Black, and Asian were 76%, 7%, and 4%, respectively.


The median baseline HbA1c was 7.2%. The mean duration of type 2 diabetes was 10.5 years and the mean BMI was 32.3 kg/m².

At baseline, 50.5% of patients had mild renal impairment (eGFR ≥ 60 but < 90 mL/min/1.73m²), 21.6% had moderate renal impairment (eGFR ≥ 30 but < 60 mL/min/1.73m²), and 1.1% of patients had severe renal impairment (eGFR < 30 mL/min/1.73m²) out of 9713 patients whose eGFR were available.

At baseline, 94.7% of patients were taking antidiabetic medication, with 10.5% of patients taking three or more antidiabetic drugs. The most common background antidiabetic drugs used at baseline were metformin (81.2%), sulfonylurea (46.0%), and insulin (23.9%). At baseline, CV disease and risk factors were managed with ACE inhibitors or angiotensin receptor blockers (81.5%), beta blockers (45.6%), calcium channel blockers (34.4%), diuretics (46.5%), statin therapy (66.1%), antithrombotic agents (58.7%), and aspirin (51.7%). During the trial, investigators were to modify antidiabetic and cardiovascular medications to achieve local standard of care treatment targets with respect to blood glucose, lipids, and blood pressure, and manage patients recovering from an acute coronary syndrome or stroke event per local treatment guidelines.

For the primary analysis, a Cox proportional hazards model was used to test for superiority. Type I error was controlled across multiple tests. TRULICITY significantly reduced the risk of first occurrence of primary composite endpoint of CV death, non-fatal MI, or non-fatal stroke (HR: 0.88, 95% CI 0.79, 0.99). Refer to Figure 6 and Table 14.

Vital status was available for 99.7% of subjects in the trial. A total of 1128 deaths were recorded during the REWIND trial. A majority of the deaths in the trial were adjudicated as CV deaths, and non-CV deaths were comparable between the treatment groups (4.4% in patients treated with TRULICITY and 5.0% in patients treated with placebo). There were 536 all-cause deaths (10.8%) in the dulaglutide group compared to 592 deaths (12.0%) in the placebo group.



Number of patients at risk

Placebo	4952	4791	4625	4437	4275	3575	742
Dulaglutide	4949	4815	4670	4521	4369	3686	741

Figure 6. KAPLAN MEIER CURVE: Time to First Occurrence of MACE in the REWIND Trial

Table 14: Treatment Effect for MACE and the Individual Components in the REWIND Trial, Median Study Observation Time of 5.4 years^a

Time to First Occurrence of:	TRULICITY N=4949	Placebo N=4952	Hazard Ratio (95% CI) ^b
Composite of non-fatal myocardial infarction, non-fatal stroke, cardiovascular death (MACE) ^d	594 (12.0%)	663 (13.4%)	0.88 (0.79, 0.99) ^c
Cardiovascular death ^{d,e}	317 (6.4%)	346 (7.0%)	0.91 (0.78, 1.06)
Non-fatal myocardial infarction ^{d,e}	205 (4.1%)	212 (4.3%)	0.96 (0.79, 1.16)
Non-fatal stroke ^{d,e}	135 (2.7%)	175 (3.5%)	0.76 (0.61, 0.95)
Fatal or non-fatal myocardial infarction ^{d,e}	223 (4.5%)	231 (4.7%)	0.96 (0.79, 1.15)
Fatal or non-fatal stroke ^{d,e}	158 (3.2%)	205 (4.1%)	0.76 (0.62, 0.94)

^a All randomized patients.

^b Cox-proportional hazards model with treatment as a factor. Type I error was controlled for the primary and secondary endpoints.

^c p=0.026 for superiority (2-sided).

^d Number and percentage of patients with events.

^e Results for components of MACE, fatal and non-fatal stroke, and fatal and non-fatal MI are listed descriptively for supportive purposes. No statistical significance should be inferred since these CIs are not adjusted for multiplicity.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

TRULICITY (dulaglutide) injection is a clear and colorless solution supplied in single-dose pens. Each TRULICITY single-dose pen is packaged in a cardboard outer carton.

Carton of 4 Single-Dose Pens

- 0.75 mg/0.5 mL solution in a single-dose pen (NDC 0002-1433-80)
- 1.5 mg/0.5 mL solution in a single-dose pen (NDC 0002-1434-80)
- 3 mg/0.5 mL solution in a single-dose pen (NDC 0002-2236-80)
- 4.5 mg/0.5 mL solution in a single-dose pen (NDC 0002-3182-80)

16.2 Storage and Handling

- Store TRULICITY in the refrigerator at 36°F to 46°F (2°C to 8°C).
- If needed, each single-dose pen can be kept at room temperature, not to exceed 86°F (30°C) for a total of 14 days.
- Do not freeze TRULICITY. Do not use TRULICITY if it has been frozen.
- Protect TRULICITY from light. Storage of TRULICITY in the original carton is recommended until time of administration.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide and Instructions for Use)

- Inform patients that TRULICITY causes benign and malignant thyroid C-cell tumors in rats and that the human relevance of this finding has not been determined. Counsel patients to report symptoms of thyroid tumors (e.g., a lump in the neck, persistent hoarseness, dysphagia, or dyspnea) to their physician [*see Boxed Warning and Warnings and Precautions (5.1)*].
- Inform patients that persistent severe abdominal pain, that may radiate to the back and which may (or may not) be accompanied by vomiting, is the hallmark symptom of acute pancreatitis. Instruct patients to discontinue TRULICITY promptly, and to contact their physician, if persistent severe abdominal pain occurs [*see Warnings and Precautions (5.2)*].
- Inform patients the risk of hypoglycemia may be increased when TRULICITY is used in combination with a medicine that can cause hypoglycemia, such as a sulfonylurea or insulin [*see Warnings and Precautions (5.3)*].
- Advise patients of the potential risk of dehydration due to gastrointestinal adverse reactions and take precautions to avoid fluid depletion. Inform patients treated with TRULICITY of the potential risk for worsening renal function and explain the associated signs and symptoms of renal impairment, as well as the possibility of dialysis as a medical intervention if renal failure occurs [*see Warnings and Precautions (5.5)*].
- Inform patients that serious hypersensitivity reactions have been reported with use of TRULICITY. Advise patients on the symptoms of hypersensitivity reactions and instruct them to stop taking TRULICITY and seek medical advice promptly if such symptoms occur [*see Warnings and Precautions (5.4)*].
- Inform patients to contact their physician if changes in vision are experienced during treatment with TRULICITY [*see Warnings and Precautions (5.7)*].
- Advise patients to inform their healthcare provider if they are pregnant or intend to become pregnant [*see Use in Specific Populations (8.1)*].
- Inform patients if a dose is missed and there are at least 3 days (72 hours) until the next scheduled dose, they should administer it as soon as possible and then resume their usual once weekly dosing schedule. If a dose is missed and the next regularly scheduled dose is due in 1 or 2 days, inform the patient to not administer the missed dose and instead resume TRULICITY with the next regularly scheduled dose [*see Dosage and Administration (2)*].
- Advise patients treated with TRULICITY of the potential risk of gastrointestinal side effects [*see Adverse Reactions (6.1)*].

Eli Lilly and Company, Indianapolis, IN 46285, USA

US License Number 1891

Copyright © 2014, 2020, Eli Lilly and Company. All rights reserved.

TRU-0012-USPI-20200903