

CLONING AND CHARACTERIZATION OF AN mRNA ENCODING AN INSULIN RECEPTOR FROM THE HORNED SCARAB BEETLE *Onthophagus nigriventris* (COLEOPTERA: SCARABAEIDAE)

Laura Corley Lavine, Laura L. Hahn, and
Ian A. Warren

*Department of Entomology, Washington State University, Pullman,
Washington*

Stephen F. Garczynski

*Yakima Agricultural Research Laboratory, USDA-ARS, Wapato,
Washington*

Ian Dworkin

*Department of Zoology, Michigan State University, East Lansing,
Michigan*

Douglas J. Emlen

*Division of Biological Sciences, University of Montana, Missoula,
Montana*

The insulin signaling pathway is the primary signaling pathway coupling growth with nutritional condition in all animals. Sensitivity to circulating levels of insulin has been shown to regulate the growth of specific traits in a dose-dependent manner in response to environmental conditions in a diversity of insect species. Alternative phenotypes in insects manifest in a

Supporting Information is available in the online issue at wileyonlinelibrary.com.

Grant sponsor: National Science Foundation; Grant numbers: IOS 0642409, IOS 0919781, IOS 0642407, IOS 0919730.

Correspondence to: Laura Corley Lavine, Department of Entomology, Washington State University, Pullman, WA 99164-6382. E-mail: lavine@wsu.edu

ARCHIVES OF INSECT BIOCHEMISTRY AND PHYSIOLOGY, Vol. 82, No. 1, 43–57 (2013)

Published online in Wiley Online Library (wileyonlinelibrary.com).

© 2012 Wiley Periodicals, Inc. DOI: 10.1002/arch.21072

variety of morphologies such as the sexually dimorphic and male dimorphic horned beetles. Large males of the sexually dimorphic dung beetle *Onthophagus nigriventris* develop a thoracic horn up to twice the length of the body whereas small males and females never develop this horn. The regulation of this dimorphism is known to be nutrition dependent for males. We focused on the insulin signaling pathway as a potential regulator of this dimorphism. We sequenced a full-length gene transcript encoding the *O. nigriventris* insulin receptor (*OnInR*), which is the receptor for circulating insulin and insulin-like peptides in animals. We show that the predicted *OnInR* protein is similar in overall amino acid identity to other insulin receptors (*InRs*) and is most closely related phylogenetically to insect *InRs*. Expression of the *OnInR* transcript was found during development of imaginal tissues in both males and females. However, expression of *OnInR* in the region where a horn would grow of small males and female was significantly higher than in the horn tissues of large males at the end of growth. This variation in *OnInR* expression between sexes and morphs indicates a role for the *InR* in polymorphic horn development. © 2012 Wiley Periodicals, Inc.

Keywords: *Onthophagus nigriventris*; insulin receptor; insulin signaling pathway; polyphenism

INTRODUCTION

Insect polyphenisms, defined as irreversible environment-specific alternative phenotypes (sensu West-Eberhard, 2003), are regulated by a complex interplay of environmental, nutritional, and hormonal factors. Variation in nutrition ultimately results in differences as extreme as the behavioral and morphological caste polymorphisms in social insects, and the remarkable sexual dimorphisms of horned beetles (Emlen, 1994; Iguchi, 1998; Moczek and Emlen, 1999; Karino et al., 2004; Wheeler et al., 2006; Emlen et al., 2007; de Azevedo and Hartfelder, 2008; Gotoh et al., 2011). The underlying genetic mechanisms responsible for these polyphenisms are still largely unknown, although studies of nutrition-dependent insect polyphenisms strongly support a key role of the insulin signaling pathway (Emlen and Allen, 2004; Emlen et al., 2006; Wheeler et al., 2006; de Azevedo and Hartfelder, 2008). The precise coordination of cell growth and proliferation in animals is regulated by the evolutionarily conserved insulin signal transduction pathway (Chan and Steiner, 2000; Claeys et al., 2002; De Meyts, 2004; Edgar, 2006; Wu and Brown, 2006; Rendell, 2008). The insulin signaling cascade in insects is initiated by the binding of insulin and insulin-like peptides to the insulin receptor (*InR*). The *InR* is a member of a large class of receptors known as receptor tyrosine kinases that have ligand-activated tyrosine kinases within their cytoplasmic domains (Hanks et al., 1988; Brogiolo et al., 2001; De Meyts, 2004). Insulin binding to the *InR* initiates a cascade of phosphorylation events resulting in the activation of glucose metabolism and protein synthesis and, importantly, cellular growth and proliferation (De Meyts, 2004; Edgar, 2006). Genetic manipulation of insulin signaling pathway components within specific imaginal discs in *Drosophila* have been shown to affect the final sizes of those traits by affecting the rate of cell proliferation (Chou et al., 1987; Potter et al., 2001; Goberdhan and Wilson, 2002, 2003; Oldham et al.,

2002; Colombani et al., 2003; Kramer et al., 2003; Puig et al., 2003; Puig and Tjian, 2005; Shingleton et al., 2005, 2008; Tang et al., 2011).

The large sword-like thoracic horn of the male dung beetle *Onthophagus nigriventris* is a striking example of an insect polyphenism. Male beetles of this species exhibit nutrition-dependent dimorphism for the thoracic horn, the length of which can extend up to twice the length of the body (Emlen et al., 2005). Males that reach a threshold body size during larval development produce a horn whereas males below the threshold body size never develop the thoracic horn. This trait is also sexually dimorphic via a nutrition-independent mechanism, as female *O. nigriventris* do not make the thoracic horn regardless of their nutritional history or body size. Physiological evidence from *Onthophagus acuminatus* and *O. taurus* (Emlen, 1997; Emlen and Nijhout, 1999, 2001) suggests that the insulin signaling pathway is either downstream or acting in concert with ecdysone and juvenile hormone to facilitate this variation in horn development and proliferation. The insulin signaling pathway has been correlated with nutrition-dependent insect polyphenisms in other species, especially in social insects (Wheeler et al., 2006; Ament et al., 2008; Azevedo and Hartfelder, 2008; Hunt et al., 2010; Wang et al., 2010; Mutti et al., 2011; Wolschin et al., 2011), has been implicated in the development of horn polymorphisms in dung beetles (Kijimoto et al., 2009; Snell-Rood et al., 2011; Snell-Rood and Moczek, 2012), and has been shown to regulate horn growth in the Asian rhinoceros beetle (Emlen et al., 2012).

The insulin signaling pathway has been suggested as a mechanism for modulating the amount of horn growth in scarab beetles in response to nutrition (Emlen et al., 2006, 2007). Emlen et al. (2006) predicted that if the insulin pathway was involved in differential horn growth, then horn discs in large males would be sensitive to circulating insulin signals while horn discs in small males and females would not. Indeed, recent studies of growth and allometry in the fruit fly *Drosophila melanogaster*, work by Tang et al. (2011), show organs differ in their size response to developmental nutrition because of differences in organ-specific specificity to circulating insulin. When faced with limited nutrition during development, male fruit fly genitalia do not respond to insulin signaling that direct the rest of the body to grow slowly (Tang et al., 2011). Based on the differential responsiveness of organs to insulin signaling, we predicted that cell proliferation in the horn discs of small males and females would be uncoupled from circulating insulin signals (Fig. 1). Indeed, we have shown differential responsiveness to insulin signaling between imaginal discs in the Asian rhinoceros beetle, *Trypoxylus dichotomus* (Emlen et al., 2012). Growing rhinoceros beetle horns are eight times more sensitive to insulin/IGF signaling than wings using RNA interference to perturb transcription of the InR (Emlen et al., 2012). Sensitivity to circulating insulin or insulin-like peptides is likely to be regulated first through the InR and then downstream by other members of the pathway (Tang et al., 2011; Emlen et al., 2012). Evidence from the development of other polymorphic animal structures such as the male-specific patterns of growth of the enlarged chelae of crustaceans and the growth of antlers in red deer (Charniaux-Cotton et al., 1966; Nagamine et al., 1980; Suttie et al., 1985, 1991; Elliot et al., 1996; Gu et al., 2007) show a critical role for insulin signaling in the development of these traits. As a first step in investigating the role of the InR in the regulation of nutrition-dependent horn polyphenisms in dung beetles, we cloned a full-length gene transcript encoding the *O. nigriventris* InR (OnInR) and examined its expression during the critical proliferative phase of horn development in males and females.




Figure 1. Diagram of the insulin signaling pathway and one model for its role in horn polyphenisms in the dung beetle *O. nigriventris*. (A) As signaling through the insulin pathway is increased overall in large males, the expression of InR decreases due to kinase-dependent inactivation of its transcriptional activator FOXO by PKB/Akt (red inhibitory bar). (B) In small males and females, insulin pathway activity is predicted to be truncated at some point downstream from the insulin receptor that would result in tissue “reprogramming” and no horn growth.

MATERIALS AND METHODS

Insects

Laboratory beetle colonies were established with *O. nigriventris* adults collected from cow and horse manure at approximately 1,000 m elevation at the Kahua Ranch, Kamuela, Hawaii 96743 in late May 2007, 2008, and 2009. Adults were kept and bred in cylindrical plastic containers 21-cm tall with a diameter of 7.5 cm containing approximately 600 ml of a sterilized soil/sand mixture and 200 ml of cow dung in an environmental chamber at 26°C with 80% humidity under a 16:8 light:dark cycle. One to two females were housed in each rearing cylinder with one male for optimal brood ball production. Brood balls were collected every 14 days from mating pairs. Brood balls were kept in 475 ml ventilated plastic deli containers on 3 cm of vermiculite using the environmental conditions described above. Beetle larvae were sexed during the third instar according to the method of Moczek and Nijhout (2002) and weights were recorded during the third instar and prepupal periods.

The transition from larva to pupa occurs in three distinct stages over a period of approximately 3 days in *O. nigriventris*. The third instar larva is the stage in which the transition to the pupal form occurs. In preparation for pupation, the third-instar larva stops feeding and initiates a purge of its gut contents. At the onset of this gut purge, the larva has entered the early prepupal period. The complete purging of the gut takes about 24 h. Other changes in the prepupa include the growth and proliferation of imaginal tissues for compound eyes, antenna, mouthparts, legs, wings, and genitalia. When the gut has completely purged, the larva has undergone many changes and has entered the late prepupal period that lasts from 24 to 48 h. Imaginal tissues thus initiate proliferation

during the early prepupal period and have completed growth and proliferation by pupation. Thus, proliferation at the early and late prepupal periods mark these as critical developmental time points when growth is regulated in these beetles and we targeted these in our experiments.

Larvae from the early and late prepupa periods were weighed and anaesthetized on ice. Under these conditions, day 2 late prepupa weight is an indicator of small (hornless) and large (horned) males and females (hornless). Larvae that will give rise to a large adult (horned male or hornless female) were at least 0.25 g on day 2 of the late prepupal period. Any individual below 0.24 g was considered to be below the threshold for a large individual and was categorized as small (hornless males and hornless females). Imaginal tissues were dissected from both large and small males and females of all sizes for RNA isolation. Imaginal tissues are easy to recognize at the early prepupal and late prepupal periods as obvious outgrowths of the larval epidermis—the horn imaginal tissues are under the thoracic dorsal region of the larval exoskeleton, wing imaginal tissues are located dorsally and laterally on the second and third thoracic segments, and the genital imaginal discs of males are found ventrally in the posterior abdominal segments.

RNA Extraction and cDNA Synthesis for Cloning

Total RNA was extracted from whole tissues dissected from late prepupal stage *O. nigriventris* stored in RNAlater[®] (Ambion, Austin, TX) using the RNeasy[®] Plus Mini kit (Qiagen, Valencia, CA) according to manufacturer's protocol for purification of total RNA from animal tissues. In order to convert RNA to cDNA, the SMART[™] cDNA amplification kit (Clontech, Mountain View, CA) and the SuperScript III First Strand Synthesis SuperMix kit and protocol (Invitrogen, Carlsbad, CA) were used. Briefly, first-strand cDNA was generated from 4 μ l total RNA primed with the CDSIII/3'PCR primer and SMART IV oligonucleotide (which is incorporated onto the 5' end of the cDNA) in a 20 μ l reaction with components supplied with the SuperScript[®] III First-Strand Synthesis SuperMix for 50 min at 50°C. Double-stranded (ds) cDNA was prepared by long distance (LD) PCR using the Advantage[™] 2 Taq Polymerase (Clontech) and protocol supplied with the SMART[™] PCR cDNA Synthesis Kit. Amplification of 2 μ l of first-strand cDNA was done by LD PCR in a 100 μ l reaction under the following conditions; initial denaturation for 3 min at 94°C, amplification for 20 sec at 94°C, 6 min at 68°C for 26 cycles followed by a 5 min 72°C incubation. The resulting ds cDNA was then purified using the High Pure PCR Product Purification Kit (Roche Applied Science, Indianapolis, IN) according to the manufacturer's protocol and eluted with 100 μ l of kit elution buffer.

The ds cDNA was normalized using duplex-specific nuclease (DSN; Axxora, LLC, San Diego, CA) according to the method of Zhulidov et al. (2004) with some modification. Denaturation and annealing was done in a total volume of 20 μ l containing 500 ng of ds cDNA in hybridization buffer (50 mM HEPES, pH 7.5, 500 mM NaCl, 0.2 mM EDTA). This mixture was heated for 5 min at 98°C to denature the cDNA and annealing was done for 4 h at 70°C. After completion of the annealing step, a 10 μ l mixture containing 3 μ l 10 \times DSN buffer (yielding a final concentration of 50 mM Tris-HCl pH 8.0, 5 mM MgCl₂, 1 mM dithiothreitol), 5.5 μ l nanopure H₂O, and 1.5 μ l DSN (0.5 Kunitz/ μ l) preheated to 70°C was added to the annealed cDNA and further incubated at 70°C for 20 min. To stop the reaction, 15 μ l of 5 mM EDTA was added. Amplification of the normalized cDNA was done by (LD) PCR and purified as described above.

Cloning and Sequencing

In order to obtain the initial fragment of the OnInR, a PCR reaction was performed in a total volume of 50 μ l using 2.0 μ l of template cDNA from *RNA Extraction and cDNA synthesis Cloning*, 31.5 μ l sterile water, 5 μ l 10 \times PCR buffer, 5 μ l 25 mM MgCl₂, 1 μ l 10 mM dNTPs, 0.25 μ l DMSO, 0.25 μ l *Taq* DNA Polymerase (Invitrogen), and 2.5 μ l of 10 μ M the following degenerate primers: InR forward 5'-CTTYGGNATGGTNTAYGARGG-3' and InR reverse 5'-CGTCATNCCRAARTCNCCRATYTT-3' (Roovers et al., 1995). The cycling conditions were as follows: a single denaturation cycle of 95°C for 10 min, followed by 35 cycles of 95°C for 30 sec, 50°C for 30 sec, and 72°C for 1 min, with a final extension step of 72°C for 10 min. An amplicon of 450 bp was detected on agarose gels and this PCR product was excised and cloned using the TOPO TA Cloning[®] kit (Invitrogen) according to manufacturer's protocol. Plasmid DNA was extracted from individual colonies using the QIAprep mini prep kit (Qiagen, Valencia, CA) and the resultant cDNA clones were sequenced at MC Laboratories (MCLab, San Francisco, CA). The cDNA sequence data were used to design sequence-specific primers for subsequent RACE reactions using Primer 3 (<http://primer3.sourceforge.net/>; Rozen and Skaletsky, 2000). For 5' and 3' RACE, the 3'-RACE CDS Primer A and the Universal Primer Mix (UPM) from the SMART[™] RACE cDNA Amplification kit were used along with the following sequence-specific primers in separate RACE reactions: OnInR 5'UTR2F 5'-GGAAGCTAGATCAACCAGAAACAATC-3', OnInR 5'UTR2R 5'-GTTTCTACAGACGGAGCATTGTTTG-3', OnInR F18F 5'-ATTTATCGGCGAAGAAGTTTGTCCATCG-3', and OnInR R22B 5'-AAGGAGGGCGGTTGCCAATAATACCAT-3' according to the manufacturer's instructions (Clontech). All PCR reactions were carried out in a Mastercycler gradient thermocycler (Eppendorf AG, Hamburg, Germany). Amplification products of the appropriate sizes were excised from agarose gels and the DNA was extracted in a GenElute[™] Minus EtBr Spin column using the manufacturer's protocol (Sigma, St. Louis, MO). Cloning and DNA sequence analysis were performed as above.

Sequence Analysis and Phylogenetic Tree Construction

Putative OnInR cDNA sequences generated above were blast searched using tBlastx (NCBI) for comparison to known InR homologues. Seqman (Lasergene software, DNASTAR[®], Inc.) was used to generate the full length OnInR gene transcript and produce a contiguous sequence from all sequencing projects. Each nucleotide was sequenced at least three times and up to 15 times. The complete open-reading frame (ORF) and translational start site of the OnInR was predicted using the ORF Finder on the NCBI website (<http://www.ncbi.nlm.nih.gov/gorf/gorf.htm>). The signal peptide was predicted using Signal 3.0 (<http://www.cbs.dtu.dk/services/SignalP/>; Nielsen et al., 1997; Bendtsen et al., 2004). Other conserved domains were predicted using the Conserved Domain Database (CDD) on the NCBI website and comparison with *D. melanogaster* InR (DInR; ACL83551) (Fernandez et al., 1995; Ruan et al., 1995), molluscan insulin-related peptide receptor (MIPR; CAA59353) from *Lymnaea stagnalis* (Roovers et al., 1995) and mosquito InR from *Aedes aegypti* (AaeIR; AAB17094; Graf et al., 1997). A protein alignment was created by the Clustal W method using MegAlign (Lasergene software, DNASTAR[®], Inc.) in order to determine similarity between the OnInR and other InRs including the human IR *Homo sapiens* (AAA59452), the mosquito *A. aegypti*, and the silkworm *Bombyx mori* (NP_001037011).

A neighbor joining analysis of OnInR with InRs from different arthropods and mammals was performed using ClustalX 2.0.12 (Larkin et al., 2007). The entire prepropeptide of each InR was entered. A diversity of animal species were chosen for the analysis including: pacific oyster (*Crassostrea gigas* CAD59674), great pond snail (*L. stagnalis* CAA59353), freshwater snail (*Biomphalaria glabrata* AAF31166), human (*H. sapiens* AAA59452), mouse (*Mus musculus* AAA39318), fruitfly (*D. melanogaster* ACL83551), mosquitoes (*A. aegypti*, AAB17094 and *Anopheles gambiae*, XP_320130), hornworm (*Manduca sexta* ACI02334), silkworm (*B. mori* NP_001037011), flour beetle (*Tribolium castaneum* EFA11583), nematode (*Caenorhabditis elegans* ACC47715), schistosome (*Schistosoma mansoni* AAV65745), and tapeworm (*Echinococcus multilocularis* CAD30260). A bootstrap analysis with 1,000 replicates was performed using MEGA 4 (Tamura et al., 2007).

Qualitative Reverse Transcription PCR and Quantitative Real-Time PCR

Qualitative reverse transcription PCR (RT-PCR) and quantitative real-time PCR (qPCR) were conducted to assess expression patterns of the OnInR transcript during the proliferative phase of prepupal development. Larvae were staged, sexed, and weighed to determine body size categories as described above. Imaginal tissues of interest included the horn area of large males, the anterior thoracic head epidermis in the region where a horn would be in small males and females, wings, legs, and male genital imaginal tissues.

Total RNA was extracted from imaginal tissues using the TRIzol[®] reagent and protocol (Invitrogen). In order to ensure no contamination from genomic DNA, the RNA was treated with DNase I according to the manufacturer's protocol (Ambion). First-strand cDNA was made using 1 µg of total RNA and the SMART[™] cDNA amplification kit and MMLV reverse transcriptase (Clontech). For all RT-PCR and qPCR experiments, single primer and no template controls were included to check for contamination. The cycling conditions for RT-PCR were as follows: 95°C for 1 min and 25 cycles of 95°C for 15 sec, 65°C for 30 sec, and 68°C for 6 min, hold at 4°C. SMART cDNAs from imaginal tissues were investigated for the presence of the OnInR transcript using sequence-specific forward and reverse primers: forward 5'-CCGAAAAGTCCAAATGGAAA-3' and reverse 5'-TCGCCGTTAGCCATTAATTC-3' and forward 5'-AACACGCAACAAACGCAGAAA-3' and reverse 5'-TAATGTCGGTTGTCCCTTGAGATAC-3'. Ribosomal 18S was used as a reference gene and primers used for *O. nigriventris* were: forward 5'-AGGGAAGACACCCTGATTCCTTCA-3' and reverse 5'-ATTCTTGGATCGTCGCAAGACGGA-3'.

For qPCR experiments, OnInR expression was assessed using primers OnInR_for: 5'-ATGAACCATCCGATTCTTGG-3' and OnInR_rev: 5'-GAGGACTTTCTTCTTGTGACG-3'. The *O. nigriventris* ribosomal 28S gene was used as a reference gene (On28S_for: 5'-CGGATCCTCCCTAACACCACATTT-3'; On28S_rev: 5'-ACAAGGATTCCCTTAGTAGC GGC-3'). All qPCR experiments were conducted using the BioRad iCycler and BioRad Sybr-Green system (Bio-Rad). Standard curves for all reference genes and for the OnInR were generated. Due to constraints on tissue sampling and tissue availability, two separate qPCR experiments were conducted. First, an experiment was conducted in large males ($n = 3$) to compare OnInR expression in horns and wings and in early prepupa and late prepupal stages. Three technical replicates and three biological replicates were performed.

The second qPCR experiment conducted was to compare small male, large male, and female horn imaginal tissues from early and late prepupa using pooled samples. In the case of large males, horn imaginal tissues from four large males at the early prepupa stage were dissected, RNA was extracted and cDNA was synthesized as described, and then the

cDNA was pooled into one sample for qPCR. This was also done for horn imaginal tissues from four large males at the late prepupal stage, thoracic epidermis in the horn region for small males, small females, and large females at the early prepupal and late prepupal stages. Four technical replicates were performed with the 28S reference gene, at the time of these experiments, no other reference genes were available. The delta Ct method, (Livak and Schmitgen, 2001) was then used to calculate the relative expression in each tissue sampled and compared using two-sample *t*-tests assuming equal variance where appropriate or else the nonparametric Wilcoxon/Mann–Whitney U test was used. This rank-based test is a nonparametric replacement test for the *t*-test and is highly conservative in comparison to equivalent rank-based tests.

RESULTS

Cloning OnInR Gene Transcript and Sequence Analysis

Degenerate primer PCR was used to amplify a 450 bp fragment from *O. nigriventris* cDNA encoding a conserved region of the InR tyrosine kinase domain (data not shown). From this fragment, sequence-specific primers were designed and used in 5' and 3'RACE reactions. Overlapping fragments generated by RACE were identified using SeqMan (Laser-gene) yielding a gene transcript of 4,757 nt. The putative OnIR transcript has a 55 nt 5'UTR followed by an open-reading frame (ORF) of 4,162 nt encoding for 1,369 amino acids and a 3'UTR of 568 nt to the polyA tail (Figs. S1–S3 and S6–S8; Genbank accession JN711464). A phylogenetic analysis of known arthropod and mammalian InRs showed that the OnInR was in fact related to other animal InRs (Fig. S4). The unrooted phylogenetic tree also fully supported the NJ tree (Figs. S4 and S5).

OnInR mRNA Expression

The OnInR transcript was first assayed for its presence in developing horn, wing, and leg tissues from large males, large females, small males, and small females using qualitative reverse-transcription PCR (RT-PCR). Male genital discs were also tested. All tissues showed the presence of the OnInR transcript (Fig. S9). Weaker bands were observed in large male horn tissue and small female wing tissue; whether this reflects an actual change would require quantitative measurements.

Quantitative real-time PCR (qPCR) was used to assess relative expression of the OnInR transcript in large males (Fig. 2). Developing horn imaginal tissues and wing imaginal tissues from the early prepupa and late prepupa stages were assayed. OnInR transcript was expressed in relatively similar levels in early prepupal and late prepupal wing imaginal tissues with no significant difference observed (Fig. 2, Wilcoxon/Mann–Whitney $\chi^2 = 2.4$, $P = 0.1213$). OnInR transcript was 1.87 times higher in late prepupa individuals in horns compared to early prepupa individuals (Fig. 2, Wilcoxon/Mann–Whitney $\chi^2 = 5.4$, $P = 0.0201$). Comparison of horns versus wings was not significantly different for early prepupa tissues (*t*-test, $t_4 = 0.205$, $P = 0.8476$) or late prepupa tissues (*t*-test, $t_{10} = 0.431$, $P = 0.6753$).

We investigated OnInR expression more closely in horn imaginal tissues in large males compared to the thoracic epidermis (corresponding region of the horn of large males) of small males and females with qPCR (Fig. 3). Comparison of OnInR transcript relative expression in small male and female anterior thoracic epidermal imaginal tissues,




Figure 2. Relative expression of OnInR transcripts in large male prepupal wing and horn imaginal tissues from two developmental stages, the early and late prepupa periods.




Figure 3. Differential expression of OnInR in horn imaginal tissues from large males, small males, and females during early and late prepupal development. Diagram of larval to adult development and graphs of the relative expression levels of OnInR during early prepupal and late prepupal development. Asterisks indicate significance of $P < 0.0001$.

to large male horn imaginal tissues during the early prepupal period was not significantly different (Fig. 3; large males to small males $t_8 = 2.315$, $P = 0.05$ and large males to females, $t_8 = 0.9425$, $P = 0.3735$). However, in the late prepupal period, the relative expression of OnInR was significantly different between small males and large males (Fig. S9; $t_8 = 9.74$,

$P < 0.0001$) and large males and females (Fig. 3; $t_8 = 8.19$, $P = 0.00011$). There were no significant differences in OnInR expression between small males and females (Fig. 3; $t_8 = 0.1474$, $P = 0.8864$). These results support what was observed in the qualitative RT-PCR assay (Fig. S9).

DISCUSSION

OnInR Protein Structure

The predicted OnInR protein is structurally similar to other known animal and insect InRs (Figs. S1–S8). Like InRs from other species, it appears that the mature OnInR protein is a tetramer composed of two α and two β subunits (Hanks et al., 1988). There are several important conserved features in the OnInR transcript that include potential insulin-binding domains, a cysteine-rich region, a tyrosine kinase domain, a furin-like site for proteolytic processing that potentially specifies the cleavage of the α from the β subunit, juxtamembrane and transmembrane domains (Figs. S6–S8). The potential α subunit contains the leucine-rich repeat domains (L1 and L2) and cysteine-rich domains (Figs. S1 and S6). Conservation of cysteine residues in the cysteine-rich domain may point to similar structural organization for the OnInR compared to other IRs (Figs. S1 and S6). Intramolecular di-sulfide bonds between these cysteines link the α subunits together and the α subunits to the β and therefore are implicated in the InR tertiary structure. The overall structure of the putative OnInR is homologous to the mammalian InR and InRs of other invertebrates (Figs. S4 and S5).

Localization of Expression

In order to determine the likelihood of the insulin signaling pathway's involvement in horn and trait growth, we investigated the distribution of OnInR mRNA in *O. nigriventris* imaginal discs during the early and late prepupal period, which encompasses the entire period of trait growth. OnInR mRNA was detected in all imaginal discs from both large and small males and females (Fig. S9). This result was expected because during the larval stages of *Drosophila*, prominent expression of InR mRNA is observed in the imaginal discs (Garofalo and Rosen, 1988). In this study, relative quantification of OnInR mRNA levels in large male horn and wing tissues during and at the end of imaginal disc growth revealed that there was essentially no difference in expression between the two stages (Fig. 2). However, when horned and hornless males were compared with each other and with females, OnInR expression differed between large males and both small males and females (Fig. 3). Specifically, by the end of the prepupal period that also marks the end of horn growth, OnInR levels in the horn tissues of large males (which had just finished growing a long horn) were significantly lower than in comparable tissues of small males and females (which did not grow horns; Fig. 3). Small males and females are distinct from large males in that they do not possess the sword-like thoracic horn.

Implications for the Mechanism of Horn Dimorphism

The insulin signaling pathway likely contributes to the nutrition-dependent modulation of growth of the exaggerated ornaments and weapons of sexual selection in animals generally (Emlen et al., 2012), and in the horns of scarab beetles specifically (Emlen et al.,

2005, 2006, 2007). This same pathway may also be part of the polyphenic mechanism shutting off horn growth in small males and females of dimorphic beetle species such as *O. nigriventris*. If the insulin signaling pathway is involved in horn dimorphism, then we predict that horn discs in large males will be sensitive to circulating insulin signals, while horn discs in small males and females should not be. Although the responsiveness of specific tissues to circulating insulin levels typically reflects the nutritional state of the individual in a dose-dependent manner, some traits are known to vary in their responsiveness to nutritional signals—through the insulin signaling pathway—affecting growth (Tang et al., 2011; Emlen et al., 2012).

As a first step toward testing the role of the insulin signaling pathway in the development of the sexually dimorphic dung beetle horn, we provide evidence that the *O. nigriventris* InR transcript is differentially expressed during horn growth and development between large horned males and hornless small males and females. Our data indicate that OnInR transcript is expressed in horn tissues during the period of horn growth, and that the levels of expression differ between individuals that have, and have not, experienced a rapid burst of proliferation in the horn imaginal tissues. Our results are therefore consistent with a role for this pathway in the polyphenic regulation of horn expression. What remains unknown is why OnInR transcript abundances were higher in the smallest males and females, rather than in the large males who were undergoing proliferative growth (Fig. 3). We suggest that this reflects an intrinsic and ancient feedback mechanism that acts to prime tissues for resuming growth when they have been exposed to starvation conditions. FOXO is a transcriptional activator for the InR (Fig. 1; Kramer et al., 2003; Puig and Tjian, 2005; Tang et al., 2011). Under starvation conditions, reduced signaling through the InR pathway can arrest cell proliferation throughout the body, but because of the feedback between FOXO and InR, starved cells begin to upregulate transcription of InR (Fig. 1). This is thought to “prime” the cells for a rapid return to full growth as soon as nutrient conditions improve and insulin-like peptides begin to circulate again (Puig and Tjian, 2005). In principle, if the OnInR pathway were interrupted at some point downstream of the OnInR in the horn tissues of small males and females, then the polyphenic mechanism of horn dimorphism would effectively mimic starvation within this tissue, activating FOXO and blocking proliferation in the horns (Fig. 1). However, activated FOXO would also signal an increase in OnInR transcription, and could generate the patterns of differential OnInR expression that we observe. While there is some evidence that the *O. nigriventris* FOXO transcript is differentially expressed in the pupal stages of male genitalia, no differences in FOXO expression was found in pupal horns between males and females (Snell-Rood and Moczek, 2012). However, because the pupal period marks the end of growth of the horn imaginal tissues in *O. nigriventris*, the role of FOXO during imaginal tissue growth and proliferation, when FOXO is predicted to have its primary effects, remains to be described in dung beetle horn polyphenisms.

Additional studies correlating receptor protein expression with transcript expression, as well as functional tests including insulin injections or receptor knockdown studies, will be needed in order to fully test this idea in the dung beetle. Comparative work in the Asian rhinoceros beetle, *T. dichotomus* (Emlen et al., 2012), suggests that the insulin signaling pathway has a major role in regulating polymorphic trait expression in dimorphic beetles. How widespread this signaling is in the development and evolution remains to be tested in scarab beetles. The results of this study of OnInR from the dung beetle are critical in elucidating the function of the insulin signaling pathway in the diversity of horned scarab beetle species specifically and its role in the control of horn growth and insect polyphenisms in general.

ACKNOWLEDGMENTS

We thank Tim Richards (Kahua Ranch, HI) for hospitality, lodging, and access to prime dung beetle habitat. We are grateful to Ralph Jennings (Pullman WA) for access to fresh, pesticide-free cow manure. Many undergraduates have helped maintain the beetle colonies including Keriann Bennett, Charlotte Seidel, Claire Walsh, Jakob Wells, and Justin Skoczylas. This work was supported by the National Science Foundation IOS 0642409 and IOS 0919781 to D.J.E. and IOS 0642407 and IOS 0919730 to L.C.L.

LITERATURE CITED

- Ament SA, Corona M, Pollock HS, Robinson GE. 2008. Insulin signaling is involved in the regulation of worker division of labor in honeybee colonies. *Proc Natl Acad Sci USA* 105:4226–4231.
- Bendtsen JD, Nielsen H, von Heijne G, Brunak S. 2004. Improved prediction of signal peptides: SignalP 3.0. *J Mol Biol* 340:783–795.
- Brogio W, Stocker H, Ikeya T, Rintelen F, Fernandez R, Hafen E. 2001. An evolutionarily conserved function of the *Drosophila* insulin receptor and insulin-like peptides in growth control. *Curr Biol* 11:213–221.
- Chan SJ, Steiner DF. 2000. Insulin through the ages: phylogeny of a growth promoting and metabolic regulatory hormone. *Am Zool* 40:213–222.
- Charniaux-Cotton H, Zerbib C, Meusy JJ. 1966. Monographi de la glande androgene des Crustaceans superieurs. *Crustaceana* 10:113–136.
- Chou CK, Dull TJ, Russell DS, Gherzi R, Leibold D, Ullrich A, Rosen OM. 1987. Human insulin receptors mutated at the ATP-binding site lack protein tyrosine kinase activity and fail to mediate postreceptor effects of insulin. *J Biol Chem* 262:1842–1847.
- Claeys I, Simonet G, Poels J, Van Loy T, Vercammen L, De Loof A, Vanden Broeck J. 2002. Insulin-related peptides and their conserved signal transduction pathway. *Peptides* 23:807–816.
- Colombani J, Raisin S, Pantalacci S, Radimerski T, Montagne J, Leopold P. 2003. A nutrient sensor mechanism controls *Drosophila* growth. *Cell* 114:739–749.
- De Azevedo SV, Hartfelder K. 2008. The insulin signaling pathway in honey bee (*Apis mellifera*) caste development—differential expression of insulin-like peptides and insulin receptors in queen and worker larvae. *J Insect Physiol* 54:1064–1071.
- De Meyts P. 2004. Insulin and its receptor: structure, function, and evolution. *Bioessays* 26:1351–1362.
- Edgar BA. 2006. How flies get their size: genetics meets physiology. *Nature Rev Gen* 7:907–916.
- Elliot JL, Oldham JM, Ahser GW, Molan PC, Bass JJ. 1996. Effect of testosterone on binding of insulin-like growth factor-I (IGF-I) and IGF-II in growing antlers of fallow deer (*Dama dama*). *Growth Regul* 6:214–221.
- Emlen DJ. 1994. Environmental control of horn length dimorphism in the beetle *Onthophagus acuminatus* (Coleoptera: Scarabaeidae). *Proc R Soc Lond B* 256:131–136.
- Emlen DJ. 1997. Alternative reproductive tactics and male-dimorphism in the horned beetle *Onthophagus acuminatus* (Coleoptera: Scarabaeidae). *Behav Ecol Sociobiol* 41:335–341.
- Emlen DJ, Allen CE. 2004. Genotype to phenotype: physiological control of trait size and scaling in insects. *Integr Comp Biol* 43:627–634.
- Emlen DJ, Nijhout HF. 1999. Hormonal control of male horn length dimorphism in the dung beetle *Onthophagus taurus* (Coleoptera: Scarabaeidae). *J Insect Physiol* 45:45–53.
- Emlen DJ, Nijhout HF. 2001. Hormonal control of male horn length dimorphism in the dung beetle *Onthophagus taurus* (Coleoptera: Scarabaeidae): a second critical period of sensitivity to juvenile hormone. *J Insect Physiol* 47:1045–1054.

- Emlen DJ, Marangelo J, Ball B, Cunningham CW. 2005. Diversity in the weapons of sexual selection: horn evolution in the beetle genus *Onthophagus* (Coleoptera: Scarabaeidae). *Evolution* 59:1060–1084.
- Emlen DJ, Szafran Q, Corley LS, Dworkin I. 2006. Insulin signaling and limb-patterning: candidate pathways for the origin and evolutionary diversification of beetle 'horns.' *Heredity* 97:179–191.
- Emlen DJ, Corley Lavine L, Ewen-Campen B. 2007. On the origin and evolutionary diversification of beetle horns. *Proc Natl Acad Sci USA* 104:S8661–S8668.
- Emlen DJ, Warren I, Johns A, Dworkin I, Corley Lavine L. 2012. A mechanism of extreme growth and reliable signaling in sexually selected ornaments and weapons. *Science* 337:860–864.
- Fernandez R, Tabarini D, Aspiasu N, Frasch M, Schlessinger J. 1995. The *Drosophila* insulin receptor homolog: a gene essential for embryonic development encodes two receptor isoforms with different signaling potential. *EMBO J* 14:3373–3384.
- Garofalo RS, Rosen OM. 1988. Tissue localization of *Drosophila melanogaster* insulin receptor transcripts during development. *Mol Cell Biol* 8:1638–1647.
- Goberdhan DCI, Wilson C. 2002. Insulin receptor-mediated organ overgrowth in *Drosophila* is not restricted by body size. *Dev Genes Evol* 212:196–202.
- Goberdhan DC, Wilson C. 2003. The functions of insulin signaling: size isn't everything, even in *Drosophila*. *Differentiation* 71:375–397.
- Gotoh H, Cornette R, Koshikawa S, Okada Y, Corley Lavine L, Emlen DJ, Miura T. 2011. Juvenile hormone regulates extreme mandible growth in the male stag beetle *Cyclommatus metallifer*. *PLoS ONE* 6:e21139.
- Graf RS, Neuenschander S, Brown MR, Ackermann U. 1997. Insulin mediated secretion of ecdysteroids from mosquito ovaries and molecular cloning of the insulin receptor homologue from ovaries of blood-fed *Aedes aegypti*. *Insect Mol Biol* 6:151–163.
- Gu LJ, Mo EY, Yang ZH, Zhu XM, Fang ZM, Sun BS, Wang CY, Bao JF, Sung CK. 2007. Expression and localization of insulin-like growth factor-I in four parts of the red deer antler. *Growth Factors* 25:264–279.
- Hanks SK, Quinn AM, Hunter T. 1988. The protein tyrosine kinase family: conserved features and deduced phylogeny of the catalytic domains. *Science* 241:42–52.
- Hunt JH, Wolschin F, Henshaw MT, Newman TC, Toth AL, Amdam Gro V. 2010. Differential gene expression and protein abundance evince ontogenetic bias toward castes in a primitively eusocial wasp. *PLoS One* 5:e10674.
- Iguchi Y. 1998. Horn dimorphism of *Allomyrina dichotoma septentrionalis* (Coleoptera: Scarabaeidae) affected by larval nutrition. *Ann Entom Soc Am* 91:845–847.
- Karino K, Seki N, Chiba M. 2004. Larval nutritional environment determines adult size in Japanese horned beetles *Allomyrina dichotoma*. *Ecol Res* 19:663–668.
- Kijimoto T, Costello J, Tang Z, Moczek AP, Andrews J. 2009. EST and microarray analysis of horn development in *Onthophagus* beetles. *BMC Genom* 10:504.
- Kramer JM, Davidge JT, Lockyer JM, Staveley BE. 2003. Expression of *Drosophila* FOXO regulates growth and can phenocopy starvation. *BMC Dev Biol.* 3:5.
- Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG. 2007. Clustal W and Clustal X version 2.0. *Bioinformatics* 23:2947–2948.
- Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* 25:402–408.
- Moczek AP, Emlen DJ. 1999. Proximate determination of male horn dimorphism in the beetle *Onthophagus taurus* (Coleoptera: Scarabaeidae). *J Evol Biol* 12:27–37.
- Moczek AP, Nijhout F. 2002. A method for sexing third instar larvae of the genus *Onthophagus* LATREILLE (Coleoptera: Scarabaeidae). *Coleop Bull* 56:270–284.

- Mutti NS, Dolezal AG, Wolschin F, Mutti JS, Gill KS, Amdam GV. 2011. IRS and TOR nutrient-signaling pathways act via juvenile hormone to influence honey bee caste fate. *J Exp Biol* 214:3977–3984.
- Nagamine C, Knight AW, Maggenti A, Paxma G. 1980. Masculinization of female *Macrobrachium rosenbergii* (de Man) (Decapoda, Palaemonidae) by androgenic gland implantation. *Gen Comp Endocrinol* 41:442–457.
- Nielsen H, Engelbrecht J, Brunak S, von Heijne G. 1997. Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites. *Protein Eng* 10:1–6.
- Oldham S, Stocker H, Laffargue M, Wittwer F, Wymann M, Hafen E. 2002. The *Drosophila* insulin/IGF receptor controls growth and size by modulating PrdInsP3 levels. *Development* 129:4103–4109.
- Potter CJ, Huang H, Xu T. 2001. *Drosophila* Tsc1 functions with Tsc2 to antagonize insulin signaling in regulating cell growth, cell proliferation, and organ size. *Cell* 105:357–368.
- Puig O, Tjian T. 2005. Transcriptional feedback control of insulin receptor by dFOXO1/FOXO1. *Genes Dev* 15:2435–2446.
- Puig O, Marr MT, Ruhf ML, Tjian R. 2003. Control of cell number by *Drosophila* FOXO: downstream and feedback regulation of the insulin receptor pathway. *Genes Dev* 17:2006–2020.
- Rendell M. 2008. Insulin: moments in history. *Drug Dev Res* 69:95–100.
- Roovers E, Vincent ME, van Kesteren E, Geraerts WPM, Planta RJ, Vreugdenhil E, van Heerikhuizen H. 1995. Characterization of a putative molluscan insulin-related peptide receptor. *Gene* 162:181–188.
- Rozen S, Skaletsky HJ. 2000. Primer3 on the WWW for general users and for biologist programmers. In: Krawetz S, Misener S, editors. *Bioinformatics methods and protocols: methods in molecular biology*. Totowa, NJ: Humana Press. p 365–386.
- Ruan Y, Chen C, Cao Y, Garofalo RS. 1995. The *Drosophila* insulin receptor contains a novel carboxyl-terminal extension likely to play an important role in signal transduction. *J Biol Chem* 270:4236–4243.
- Shingleton AW, Das J, Vinicius L, Stern DL. 2005. The temporal requirements for insulin signaling during development in *Drosophila*. *PLoS Biology* 3:1607.
- Shingleton AW, Mirth CK, Bates PW. 2008. Developmental model of static allometry in holometabolous insects. *Proc R Soc Lond B* 275:1878–1885.
- Snell-Rood EC, Moczek AP. 2012. Insulin signaling as a mechanism underlying developmental plasticity: the role of FOXO in a nutritional polyphenism. *PLoS One* 7:e34857.
- Snell-Rood EC, Cash A, Han MV, Kijimoto T, Andrews A, Moczek AP. 2011. Developmental decoupling of alternative phenotypes: insights from the transcriptomes of horn-polyphenic beetles. *Evolution* 65:231–245.
- Suttie JM, Gluckman PD, Butler JH, Fennessy PF, Corson ID, Laas FJ. 1985. Insulin-like growth factor 1 (IGF-1) antler-stimulating hormone? *Endocrinology* 116:846–848.
- Suttie JM, Corson ID, Gluckman PD, Fennessy PF. 1991. Insulin-like growth factor 1, growth and body composition of red deer stags. *Anim Prod* 53:237–242.
- Tamura K, Dudley J, Nei M, Kumar S. 2007. MEGA4: molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mole Biol Evol* 24:1596–1599.
- Tang HY, Smith-Caldas MSB, Driscoll MV, Salhadar S, Shingleton AW. 2011. FOXO regulates organ-specific phenotypic plasticity in *Drosophila*. *PLoS Genet* 7:e1002373.
- Wang Y, Mutti NS, Ihle KE, Siegel A, Dolezal AG, Kaftanoglu O, Amdam GV. 2010. Down-regulation of honey bee IRS gene biases behavior toward food rich in protein. *PLoS Genet*. 6, e1000896.
- West-Eberhard MJ. 2003. *Developmental plasticity and evolution*. Oxford, United Kingdom: Oxford University Press, Inc.
- Wheeler DE, Buck N, Evans JD. 2006. Expression of the insulin pathway genes during the period of caste determination in the honey bee, *Apis mellifera*. *Insect Mol Biol* 15:597–602.

- Wolschin F, Mutti NS, Amdam Gro V. 2011. Insulin receptor substrate influences female caste development in honeybees. *Biol Lett* 7:112–115.
- Wu Q, Brown MR. 2006. Signaling and function of insulin-like peptides in insects. *Ann Rev Entom* 51:1–24.
- Zhulidov AV, Bogdanova EA, Shcheglov AS, Vagner LL, Khaspekov GL, Kozhemyako VB, Matz MV, Meleshkevitch E, Moroz LL, Lukyanov SA, Shagin DA. 2004. Simple cDNA normalization using Kamchatka crab duplex-specific nuclease. *Nucleic Acids Res* 32:e37.

Supplementary Materials

1 M A
 1 GATGCATGCTCTGAGCGCAAAGCGGGCCTCTGCTTTGCCGGATAAGTCACCCCATGGCA
 3 S K S I C G L A D T G R L C S W L L V A
 61 TCGAAGTCGATTTGCGGACTTGCTGACACGGGCCGGCTCTGTAGTTGGTTGCTGGTTGCG
 23 W V V F G F Y C N L P V T H A G Y I T E
 121 TGGGTCGTTTTTGGATTTTACTGCAACTTGCCGGTTACCCACGCTGGTTACATTACCGAA
 43 Y N G F T P Y N K T S G I C K S V D I R
 181 TACAACGGATTTACTCCATATAATAAAACATCAGGCATATGCAAGAGTGTGGACATTCGA
 63 N S G E M F E N L R G C R V V E G Y I Q
 241 AATTCAGGGGAGATGTTTCGAGAACTTGCGAGGTTGCAGAGTGGTTGAGGGTTACATCCAA
 83 I L L F D H E N K S T F E N I S F P E L
 301 ATCCTTCTTTTCGACCATGAAAATAAATCGACGTTTGAAAACATCAGCTTTCCCGAATTG
 103 R E I T G H L L L Y R V N G L T S L G K
 361 CGGAAATTACTGGTCATCTACTCTTGTATAGGGTGAACGGATTGACTAGTTTGGGTAAA
 123 L F P N L A V I R G Q Q P L L L G Y A L
 421 CTTTTTCCGAAGTTGGCCGTGATAAGAGGTCAACAGCCGTTGTTGTTGGGGTACGCGTTG
 143 I I F E M P S L Q E I G L Y S L T D I M
 481 ATCATCTTTGAAATGCCTAGTCTGCAAGAGATCGGTCTCTATTTCGTTAACGGACATTATG
 163 N G G I R I D K N P Q L C F A N T I N W
 541 AACGGTGGGATTTCGGATTGATAAAAATCCTCAATTGTGCTTCGCTAATACGATCAACTGG
 183 N R I M H K K N K D A E L F I R S S K K
 601 AATAGGATTATGCACAAGAAAAATAAAGATGCCGAAGTGTATAAGGAGTAGTAAAAAA
 203 E N E C P M C P R M D N G K P L C E Y C
 661 GAAAATGAGTGTCCGATGTGTCCGAGGATGGATAACGGAAAACCTTTGTGCGAGTATTGT
 223 W S S K H C Q K K C K S N C T S C N D L
 721 TGGAGTAGCAAACACTGCCAGAAAAAATGTAAGAGTAATTGTACGTCGTGCAACGATTTG
 243 G E C C N N T C I G G C S N N D P K Q C
 781 GGGGAATGTTGTAATAATACGTGTATCGGCGGGTGCTCGAACAATGACCCCAAACAATGC
 263 S V C R N I T F Y T E F N K T V C L D K
 841 TCCGTCTGTAGAAACATTACCTTTTACACGGAATTTAATAAAACAGTGTGTCTAGACAAG
 383 C P E G L F K Y L D R R C V T R E E C I
 901 TGCCCAGAGGGTTTGTTTAAATATTTAGATAGGAGATGCGTAACTAGAGAAGAATGTATT
 303 N S P K H I E Y T T S K Q E Y E F K T F
 961 AACTCGCCAAAACACATAGAATACACCACCAGCAAGCAAGAGTACGAATTCAAAACCTTTT
 323 N N T S C V Y K C P P K Y T D D P K K K
 1021 AATAACACCAGTTGTGTCTACAAATGTCCACCGAAATACACCGACGATCCGAAGAAAAAA
 343 A C V S C D S R C K K V C P G I S V N N

1081 GCGTGC~~TTTTCTTGCGACTCTAGATGCAAAAAGGTTTGTCCAGGAATAAGTGTTAACAAT~~
 363 *I E T A E S L K E* **C** *T H I T G S I E I E*
 1141 ~~ATTGAAACAGCAGAATCCCTAAAAGAGTGC~~**ACTCAT**~~ATTACTGGATCTATAGAAATCGAA~~
 383 *V S G K N I V P A L N A N L N M I E E I*
 1201 ~~GTTAGTGGAAAAAATATTGTACCTGCATTAACGCAA~~**ACTTAA**~~CATGATTGAGGAAATA~~
 403 *D G* **Y** *L K I V R S* **Y** *P L I S L S F F K S*
 1261 ~~GACGGTTATTTAAAAATAGTTAGATCATATCCGTTGATATCACTAAGTTTTTTCAA~~**AA**~~AGT~~
 423 *L K V I H G N A F E N G K* **Y** *S F I V L D*
 1321 ~~CTCAAGGTGATTCACGGAAATGCTTTTGAAAACGGAAAGTATTCGTTTATCGTTTTGGAC~~
 443 *N Q N L Q N F W D W E N R T L Q I N H G*
 1381 ~~AATCAGAATTTACAAA~~**ACTTCTGGGATTGGGAAAACCGAACGTTGCAAATTAACCACGGA**
 463 *K L F F H F N P K L* **C** **Y** *K L I K Q L A E*
 1441 ~~AAATTGTTCTTCCACTTTAACCCAAAGTTGTGTTACAAATTAATCAAACA~~**ACTCGCCGAA**
 483 *K A N I T K F D D W D V A E N S N G D K*
 1501 ~~AAAGCTAACATTACTAAATTCGACGATTGGGATGTGGCGGAAAATTC~~**CAACGGCGATAAA**

Supplementary Figure 1. OnInR α subunit. Signal peptide (grey shade), leucine-rich repeats (italic) and cysteine-rich domain (bold). Conserved tyr are highlighted in green and conserved cys in yellow.

503 T A C E I K D L I I E I Q S K S S T S V
 1561 ACCGCTTGTGAAATAAAAGACCTAATTATAGAAATTCAGTCGAAATCGAGCACATCAGTT

523 Q L Y W V P F N I T D S R K L L S Y Q I
 1621 CAACCTTTATTGGGTTCCGTTTAATATCACCGATTCCCGGAAATTACTTTCTTATCAAATT

543 Y H I E A P V Q N F S I F D S R D A C G
 1681 TATCATATTGAAGCACCCGTACAAAATTTCTCCATATTTGATAGTAGAGATGCTTGTGGG

563 G D G W S I Q D V P I P Y D H H N V S K
 1741 GGGGATGGATGGTCAATACAAGATGTTCCAATCCCTTATGATCACCACAACGTATCAAAA

583 I E A T L K D L K A Y T Q Y A F F V K T
 1801 ATCGAAGCAACCCTAAAAGATTTAAAAGCATACACGCAATACGCTTTTTTCGTCAAAACT

603 Y T I A Q E K N G G L S A L K Y F R T N
 1861 TACACCATAGCACAAAGAAAAAATGGCGGATTAAGTGCTCTCAAGTATTTTCAGAACAAC

623 P G Q P S S P T F L N L R A N S S S S I
 1921 CCTGGCCAACCTTCATCGCCGACTTTCTTAAATTTACGAGCAAATAGCAGTAGCAGTATA

643 E V D W S P P K S P N G N L T H Y K V I
 1981 GAAGTCGACTGGTCACCGCCGAAAAGTCCAATGGAAATTTAACTCATTATAAAGTAATC

663 V T L K E T Q D E L E Y S N S C S V R S
 2041 GTTACTCTGAAGGAAACTCAAGATGAATTAGAATATAGTAACTCGTGCTCTGTAAGGTCT

683 S Q T T P K T S D P A K P T D S P P S K
 2101 TCCCAGACAACCTCTAAAACCTTCAGACCCGGCAAAACCAACAGATTCTCCTCCATCAAAA

703 N D T C D C T T D M L R E S N A L A N K
 2161 AACGACACCTGCGATTGCACCACCGATATGCTTCGCGAATCAAATGCGCTCGCTAACAAA

723 E A E T E S R I N F E N E L Q N R V Y I
 2221 GAAGCGGAAACGGAATCTCGAATTAATTTTTGAAAATGAACTACAAAATCGGGTTTATATT

743 K R V A P P P Q P S M T L S R K K R Q M
 2281 AAAAGAGTTGCCCCACCACCACAACCTTCGATGACTCTTAGCAGAAAAAAAAGACAAATG

763 F P S H N D S N V S D S Y V G N K T Y V
 2341 TTTCTTCACATAATGATTCAAACGTTTCCGATTCATACGTAGGTAATAAAACTTACGTT

783 A T Y I V T G T K F Y I T N L Q H H T S
 2401 GCCACTTATATTGTGACTGGTACTAAATTCTATATAACAAATTTACAACATCATACGAGT

803 Y E I S V Q A C R K T E P D K G D F Q P
 2461 TATGAAATATCAGTACAAGCATGTAGGAAAACGGAACCTGACAAAGCGATTTTTCAACCT

823 N C S D S T S K E I R T E S K H D A D N
 2521 AATTGCAGCGATAGTACTTCAAAGAAATTCGAACGGAAGCAAACATGATGCGGATAAT

843 I A S N I S I V S V T E N S I T I S W K
 2581 ATTGCTTCAAACATATCGATAGTTTTCGGTTACGGAAAATTCATAACAATATCTTGAAA

863 P P I N P N G R I R H Y F I D V S K S G
 2641 CCTCCGATAAATCCCAACGGACGTATCCGACATTATTTTCATAGATGTTTCAAATCAGGA

883 T P I P E R C I D N Q D F K N S S Y K Y
 2701 ACTCCAATCCCTGAACGCTGCATTGACAACCAGGATTTTAAAAATAGCAGCTACAAATAT

 903 T I E P L S P G N Y T I R V Q A I S D A
 2761 ACTATTGAACCGTTATCTCCTGGGAATTATACGATACGAGTTCAAGCGATATCAGATGCA

 923 N T G G A F S L P V S V Y I D E P S D S
 2821 AATACTGGCGGTGCTTTTTTCATTACCGGTTTCTGTCTATATCGATGAACCATCCGATTCT

Supplementary Figure 2. Fibronectin type III domains and processing site. Interdomain contacts (grey shade), FnIII-1 (*italic*) and 3 (underline) domains, cleavage site (fuchsia shade), conserved cysteines (yellow highlight, bold-disulfide bonds) and tyrosines (green highlight).

943 W N I I A G I I I L V L I V V G I I V Y
 2881 TGGAATATTATAGCGGGAATTATTATTCTTGTTTTAATCGTTGTGGGTATTATTGTATAC

 963 I L F K K R Q Q E E S P R M N P S I N P
 2941 ATACTATTTAAAAACGTCAACAAGAAGAAAGTCCCTCGGATGAATCCGTCAATAAATCCC

 983 E Y M Y V P D D W E I P R K K I E L H K
 3001 GAATATATGTATGTTCCGGACGATTGGGAAATTCCACGGAAAAAATTGAATTACACAAA

 1003 E L G Q G S F G M V Y E G I V R D I K G
 3061 GAGTTGGCCAAGGAAGTTTTGGCATGTTTATGAAGGAATTGTACGTGATATTAAGGA

 1023 K A T I R C A I K T V N E H A T N A E R
 3121 AAAGCGACGATTCGTTGTGCTATTA AACCGTTAACGAACACGCAACAAACGCAGAAAAGG

 1043 L N F L K E A S V M K A F D T A H V I K
 3181 TTAAATTTCTTAAAAGAAGCTTCGGTTATGAAAGCATTGATACAGCGCACGTGATAAAA

 1063 L L G V V S Q G Q P T L V V M E L M A N
 3241 CTATTAGGAGTTGTATCTCAAGGACAACCGACATTAGTTGTAATGGAATTAATGGCTAAC

 1083 G D L K S Y L R L H R P D S D G Y T H G
 3301 GGCGATTAAAATCTTATCTTAGGTTACACAGACCAGATTCAGATGGTTACACTCATGGT

 1103 I I G Q P P S L R R I L Q M A I E I A D
 3361 ATTATTGGGCAACCGCCCTCCTTACGAAGGATATTACAAATGGCGATAGAAATCGCTGAC

 1123 G M A Y L S A K K F V H R D L A A R N C
 3421 GGTATGGCTTATTTATCGGCGAAGAAGTTTGTCCATCGTGATTTAGCGGCGAGAAATTGT

 1143 M V A E D L T V K I G D F G M T R D I Y
 3481 ATGGTAGCGGAAGATTTAACTGTTAAAATAGGCGATTTTGGTATGACTAGAGATATTTAT

 1163 E T D Y Y R K G T K G L L P V R W M A P
 3541 GAAACGGATTATTATCGAAAAGGAACAAAGGGATTGTTACCAGTTAGGTGGATGGCACCG

 1183 E S L K D G V F S S S S D I W S Y G V V
 3601 GAAAGTCTTAAAGACGGAGTTTTTTCAAGTAGTAGTGATATTTGGAGTTATGGCGTGGTA

 1203 L W E M A T L A S Q P Y Q G L S N D Q V
 3661 TTGTGGGAGATGGCTACTTTAGCATCACAACTTACCAAGGATTATCGAACGATCAAGTC

 1223 L R Y V I D S G I M E R P E N C P D K L
 3721 CTAAGGTATGTTATTGATAGCGGTATTATGGAAAGACCAGAAAATTGTCCGGATAAACTC

 1243 Y H L M R Q C W K Y K Y A E R P S F L D
 3781 TATCATTTGATGCGACAATGTTGGAAGTACAAATACGCGGAAAGACCGAGTTTTCTCGAT

 1263 L V S M L L E D S S P Q F S R V S F Y H
 3841 TTAGTCTCGATGTTGTTGGAGGATTCGAGCCCGAATTTAGCAGGGTTTTCGTTTTATCAC

 1283 S A A G M E A R S T R N N Q M I I Q D D
 3901 AGCGCGGCTGGAATGGAAGCTAGATCAACCAGAAACAATCAAATGATCATTCAAGACGAC

 1303 V M M P L R L T K E L E E I Y P L N S C
 3961 GTAATGATGCCGTTAAGACTGACCAAAGAAGACTAGAGGAGATTTATCCGTTAAATAGTTGC

 1323 A D D S E V E D D L V G S P H M E F S S



```

4021 GCTGACGATTCGGAAGTCGAAGACGATTTGGTTGGAAGCCCGCATATGGAATTCTCCAGC
1343 Y P K V Q K D T T T G T A N G Y V G T N
4081 TATCCAAAAGTGCAAAAGGACACAACACTACAGGTAAGTGGATATGTTGGAACGAAC


1363 P S N G T R A -
4141 CCATCGAATGGTACTAGAGCGTAAAAAAGGAAAAAAAAAAAAACAAATGGGAAATTCTCGA
4201 AATAGTAGTTCTACCAAAGATAGTGATAAAAGAATAAAAAATAAACTAAAAAAGTGTAT
4261 CGATATGAAATAATGCAGAACCGATCGTATACTCTACTTACCACTCATTCTGAATTGTATT
4321 CCATTTTATTTTTTTAAAAGGGAATTTAAAAAATAGTTCATCTTGCCCAAGTACCGAAC
4381 GACGTGCGAGTTTTAGCCATTGCCTTATGTAAAATTGATTAATTGATTGCGTGCCCTAAG
4441 CGCGCTGCACCCCCTCCCTCAGACCTATGGTCATCCTCTATTAGTTTTGGGTTTCGTCGC
4501 CCGCAATCTCTACAGCATAAGCCTAAGTTCCTGATATGACCATAATTTAAATCCGATATT
4561 TTATAAATTAATAAAAAATCAAGCATAGAATGGTCGTTAAAAACGTAAGTGAATTAATT
4621 TTTATCATACTACTACTGTTATTTATATTTTTAATTCGTTATATTTTGTACCATTGAGA
4681 AAATTGTACATTCTGAACAATTTACTTTTTATATAAAATTATTGTAGGATATGAAAAAAA
4741 AAAAAAAAAAAAAAAAAA

```

Supplementary Figure 3. OnInR β subunit. Transmembrane domain (TMD; bold), juxtamembrane domain (JMD; italic), beginning and end of the tyrosine kinase domain (TKD; grey shade), ATP-binding site (fuchsia shade), protein kinase and protein tyrosine kinase motifs (blue and red shade).



Supplementary Figure 4. Phylogenetic tree of several vertebrate and invertebrate IRs. Entire prepropeptide sequences were used. The tree was generated with ClustalW alignment and neighbor-joining tree with bootstrap analysis. Bootstrap values are shown at the nodes of the branches and represent the percentage of times that grouping was supported. Representatives from the molluscs were used as an outgroup.



Supplementary Figure 5. Comparative InR unrooted phylogenetic tree. Entire prepropeptide sequences were used. An unrooted phylogenetic tree was created using MEGA 4 (Tamara et al., 2007). Support at each internal node was assessed using 1000 bootstrap samplings and the tree was visualized using Tree View (Page 1996). Branch lengths are drawn to scale.

Supplementary Reference:

Page, R. D. M., 1996. TREEVIEW: An application to display phylogenetic trees on personal computers. *Comp. Applic. Biosci.* 12, 357-358.

Majority	-----XGXVRRACXLLLVAAAXLLXAXAXTYAGX-X-----X-----GVCXSMDIRNXPX	
OnInR	-----MASKSICGLADTGR LCSWLLVAVVVF GFCNLVPTHAGYIT-----EYNGFTPYNKTS GICKSVDIRNSGE	66
DInR	PGISLLLFILLANTLAIQAVVLPAHQQLLHNDIADGLDKTALSVSGTQTRWRPSESNTMRLSQNVKPKCSMDIRNMVS	349
HIR	-----MGTGGRRGAAAAPLLVAVAALLGAAGHLYPG-----EVC PGMDIRNNLT	45
AaeInR	-----MALSGQNMHNLG-----VCGSV DVRNSPA	24
BomInR	-----MLEVRRGVMRRACDR LGWPAVACALLIACARTYAGTEV-----VP-----ARGVCP SMDLRNPNP	55
Majority	HXXRLEXRVI EGFLLQILLXDKAXPSXFENXSF PKLXEITXYLLYRVNGLXSLGXLFPNLAVIRGXQ-LFXXYALVIFE	
OnInR	MFENLRGCRVVEGYIQILLFDHENKSTFENISFPELREITGHLLYRVNGLTSLGKLPNLAVIRGQQPLLLGYALIIFE	146
DInR	HFNLQENCTVIEGFLLIDLINDASP---LNR SFPKL TEVTDYIIYRV TGLHSLSKI FPNLSVIRGNK-LFDGYALVVYS	425
HIR	RLHELENC SVIEGHLQILLMFKTRPEDFRDLSFPKLIMITDYLLLFRVYGLESLKDLFPNLTVIRGSR-LFFNYALVIFE	124
AaeInR	HLDR LKDCVVEGFVHILLIDKYIDSS FENYSFP LLTEITEYLLLFRV NGLKSLRRLFPNLAVIRGDA-LVGDYAMVIYE	103
BomInR	QIRRLGCRVIEGQLSIVLMERATNKIFDNMSFPQLREVTGYIL IYKTKGLRNLGDLFPNLPVVRGMQ-LFKDFALVIFD	134
Majority	XXHLEELGLYS LMDITRGGVRIEKNPKLCYANTIDWSR IXXX--EDAEIVXRSNXKENECPCPXXX-X-X---XKXXC	
OnInR	MPSLQEIGLYSLTDIMNGGIRIDKNPQLCFANTINWNRIMHKKNKDAELFIRSSKKENECPCPRMDN-----G	215
DInR	NFDLMDLGLHKLRSITRGGVRIEKNHKL CYDRTIDWLEILAEN-ESQLVVL TENGKEKESLSKCPGEIRIEEGHDNTAI	504
HIR	MVHLKELGLYNLMNITRGSVRIEKNNELCYLATIDWSRILDSV-EDNHIVLNKDDNEECGDICPGTAK-----GKTNC	196
AaeInR	LMHIEEIGLISLMDITRGGVRIEKNPKLCFANTIDWKAMTVP---GTNNYIKDNQKDNVCPICPAESTAVMLPNGSKQKC	180
BomInR	NEHLEALGLNSLMKIERGGVRIQHNDRLCYTNTIDWSRITR---DDAEIIVRSNYDTRLCGLCPNAQSRVEE--DHRQSL	209
Majority	PAXPX---KXXXREL CWSKHCQKXCPSXCXS-GCXDEGTCCNSE-CLGGCSXP--DPSSCSVCRNFSFDX---X-XCX	
OnInR	-----K-PLCEYCWSSKHCQKKCKSNCTS--CN DLGECNNT-CIGGCSNN--DPKQCSVCRNITFYTEFNKTVCL	280
DInR	EGELNASQLHNNRRLCWSNKLQTKCPEKCRN-NCIDEHTCCSQD-CLGGCVIDKNGNESCI SCRNVSFNN-----ICM	577
HIR	PATVI---NGQFVERCWTHSHCQKVCPTICKSHGCTA EGLCCHSE-CLGNCSQPD-DPTKCVACRNFYLDG-----RCV	265
AaeInR	PAAPVRGGNKDHKRTL CWNANHCQTICPPECPK-ACSKTGVCCDAESCLGGCNLP--NTSSCSVCRHLSIDPAG-KRQCV	256
BomInR	PQCPA---DTKGKLLCWDEKHCQKICPSACGDHGCMSNGTCCNSA-CLGGCDGP--LASNCFVCKKFSFEYGS-EMTCM	281

Supplementary Figure 6. Insulin receptor protein alignment of the L1 and L2 domains and the cysteine-rich region. Abbreviations: *O. niriventr* insulin receptor (OnInR), *Drosophila* insulin receptor (DIR), human insulin receptor (HIR), *A. aegypti* insulin receptor (AaeInR), *B. mori* insulin receptor (Bom InR).

Majority	XXCPPGYKXX-RRCVTRXECIXXXKPIXNSXXXXXXXF-----XXXNGSCXXECPSGYTMXXX---RXCXPCX -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+	
OnInR	DKCEPLFKYLDRRCVTRREECINSPKHIEYTTSKQEYEFK-----TFNNTSCVYKCPPKYTDDPKK---KACVSCP	348
DInR	DSCPKEYYQFD-SRCVTANECITLTKFETNSVYSG-----IPYNGQCITHCPTGYQKSENK---RMCEPCD	639
HIR	ETCPPPYHFQDWRCVNFSSFCQDLHHCKNSRRQGCHQY-----VIHNNKCIPECPSGYTMSSN---LLCTPCL	332
AaeInR	AKCPPNTFKYH-TRCVTRDECYAMKKPISLDSNPDL PDQP-----FIPHNGSCLMECPLDHELITELNKTRWCRKCS	327
BomInR	ESCPGGTYELS-RRCVTEQECRSMSPPPQVGAADSTYFRFAIKVPAYKTFNSSRCLFMCPSGY-MEVGN---ETNATCQ	356
Majority	G-PCPKVCXGSE-----IDSSEXAXELXGCTXITG--SLEIXRXR--GXNIVXELEANLGXIEEIXGYLKVVRSYPLVSL -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+	
OnInR	S-RCKKVCPGIS-----VNNIETAESLKECTHITG--SIEIEVSG---KNIVPALNANLNMIEEIDGYLKIVRSYPLISL	417
DInR	GGKCKECS-SG----LIDSLERAREFHGCTIITGTEPLTISIKRESGAHVMDPKLCVGLAAVHKIQSSLMVHLTGYLKSLSL	714
HIR	G-PCPKVCHLLEGEK-TIDSVTSAQELRGCTVING--SLIINIRG--GNNLAAELEANLGLIEEISGYLKIRRSYALVSL	406
AaeInR	G-TCPKRCEGSN-----IDNIQSAQLKGCIEIDG--SLEIQLRSRGGENIVKELENFLSSITEIKGYLKVVRSYPLLSL	399
BomInR	--PCPRSGCVRECLGGKIDSVATAEQFRGCTHIK--KMEFTLRSS-GGKTLSSLLESALGEIREIHGCLQVTRSYPLVSL	431
Majority	SFFKLLXXIHGNP-XEXGXYSLYVLDNQNQLQELWDWXX-TXTIXXGKLEFFHFNPKLCYSXIKQLXEMX--KTXFEXXDV -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+	
OnInR	SFFKSLKVIHGNA-FENGKYSFIVLDNQNQLNFWDWENRTLQINHGKLEFFHFNPKLCYKLIKQLAEKAN--ITKFDDWDV	494
DInR	KFFQSLTEISGDPPMDADKYALYVLDNRDLDELWGPNQ-TVFIRKGGVFFHFNPKLCVSTINQLLPLASKPKFFEKSD	793
HIR	SFFRKLRLIRGET-LEIGNYSFYALDNQNLRLQLDWDSKHNLTQTGKLEFFHYNPKLCLSEIHKMEEVSGT-KGRQERNDI	484
AaeInR	GFLKLLKIIHGKG-NKVSNSSLYVVENQNQLQELFDHN---VTIGEGKLEFFFNPMCLTDRKAVKKYNP--GIEIENESQ	473
BomInR	MFLKLLRKIIANP-SEDKGEALHIINNENLELLWDWSTYGPIEIIIGSCYIHSNPKLCYTQLMPLMMSYP-KTNFTELEV	509
Majority	AEDSNGDXASCEIXVLKXSLQSXSXSXXLXWPXXXX-DXRKLLGYXIYXXEAP-YQNVXFFDGRDACGXDGWVXD-- -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+	
OnInR	AENSNGDKTACEIKDLIEIQSKSSTSVQLYWVFNIT-DSRKLLSYQIYHIEAP-VQNFISIFDSRDACGGDGSIQD--	570
DInR	GADSNRGSCTAVLNVTLQSVGANSASLNVTTKVEIGEPQKPSNATIVFKDPRAFIGFVYHMIDPYGTQLKAVTIH-	872
HIR	ALKTNGDKASCENELLKFSYIRTSFDKILLRWEPYWPP-DFRDLLGFMLFYKEAP-YQNVTEFDGQDACSNSWTVVD--	560
AaeInR	LESNNGDRAACSITELTSLKSIGSETAI IQWAPFTELSDARMLLGYVIYYIEAP-YANVTFFDGRDACNTEGWRLDD--	550
BomInR	SQDSNGYQASCLPDVLLKLSVSELTQIAVVLTDWVYCPD-DMRKLGLYSLYFMAT--EKNVSLYDQRDAC-SDIWKVLDPK	585
Majority	--XXXISNXNXSXEHAG-----XLTLQLKPYTQYAYYVKTYTXXSEX--GAQSDIKYFRTNPGQPS -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+	
OnInR	-VPIPYDHHNVSKIEAT-----LKDLKAYTQYAFFVKTYTIAQEK--GGLSALKYFRTNPGQPS	627
DInR	---AMIAGKVSSPEKSG-----VMVLSNLIPTYNYSYVRTMAISSELT---NAESDVKNFRTNPGRPS	930
HIR	IDPPLRSNDPKSQNHG-----WLMRGLKPWTQYAFVKTIVTFSDERRTYGAKSDIYVQTDATNPS	623
AaeInR	-----ISDFNMDKETTK-----ILTQLKPYTQYAYYVKTYTLGSEGL---GGQSKIKYFTTAPGTPS	604
BomInR	IEETRKVNRNAKPSHARNKTEVIFNPCADMQPLFYPTQLRPYTRYAAYVKTYTTIQDRK---GAQSSIIYFRTLPMQPS	662
Majority	VPTDVXXANSSSSIXVXWSPXXPNNGXTHYXVFVELNAXDRE---XXNXDYCXXPXXLASXTXXPPTXPKPXXSPXX -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+	
OnInR	SPTFLNLRANSSSSIEVDWSPKSPNGNLTHYKVIVTLKETQDE-----LEYSNSCSVRSSQTTPKTSDBAKPTDSPPS	701
DInR	---KVVATAISDSKINVTWYLDNLMLTRYFIKAKLINRPTRN---NNRDYCTEPLVKAMENLDPATTPTKISDPLA	1003
HIR	VPLDPISVSNSSSQIILKWKPPSDPNGNITHYLVFWERQAEDSE---LFELDYCLKGLKLPRTWSPPFESDSQKHNQS	700
AaeInR	VVRDVEVSVN-KNMLTVKWLPPLKMNGLKEYEVFIELNADDNEQ--LMLRDYCED-DKLRDIPVETPTSAPPKTSICT	680
BomInR	PPTAVTVELLTPHSIFIKWSPPTMPNGTIILYHVEVQANSYNRPQILAGNINYCGNPSILANMIKASLEEAPETPEEKIT	742
Majority	---KXGXCCCEXPXXXSXLTXDEEEXEXXINFENELQNXVYXKX-X-----VNPXPXXSXRPS -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+	


		-----+-----+-----+-----+-----+-----+-----+-----+							
OnInR	---	KNDTCDCTDMLRESNALANKEAETESRINFENELQNRVYIKR-----VAPPPQPSMTLSR	757						
DInR	----	GDCKCVEGSKKTS-SQEYDDRKVVQAGMEFENALQNFIFVPNIRKSKNGSSDKSDGAEGAALDSNAIPNGGATNPS	1077						
HIR	-	EYEDSAGECCSPKTDSQLKEELESSFRKT-FEDYLHNVVFPVPR-K-----TSSGTGAEDPRPS	758						
AaeInR	---	ADQCRNYCKAPTSGGSTGTIDVTDKENQITFEDQLHNVYVIKN-----PLLRDKTTRRK	734						
BomInR	G	VDKNGVCACKEEAKPSLRFSTRAEEERVDSINFENELQNVYRKSER-----VNPKIHKSIDGTS	803						
Majority	RKRRXX-----	XXXXXXXXTEXTFP--XX----NXSNXSDEEYXXXXEXLVVXTTFVISNLRHFTX							
OnInR	KKRQ-----	MFP--SH---NDSNVSDSYVGNKYVATYIVTGTGFYITNLQHHTS	802						
DInR	RRRRDVALEPELDDVEGSVLLRHVRSITDDTDAFFEKDDENTYKDEEDLSSNKQFYEVFAKELPPNQTHFVFEKLRHFTR		1157						
HIR	RKRRSLGDV-----	GNVTVAVPTVAAFP-----NTSSTSVPTSPEEHRPFKEVVNKESLVISGLRHFTG	817						
AaeInR	RSTN-----	LLFPNNTENKNDTTD----RRTEKVKDEPYQYIFNATNETSITFPLSYFNHYSL	790						
BomInR	KVKRSVDHS-----	LNSMLVKLTENNSPRPGL----NYSNITDDEGYVKSLEYELDGSTRTLTVKNMRHFTW	866						
Majority	YXIXXQACRXXXPXEXX-X----	FEXXCXSVAYXXFRTXKXKXADDIVSNXXVEIEXANXT--XXTVSWK-XPLNPNGX							
		+-----+-----+-----+-----+-----+-----+-----+							
	970	980	990	1000	1010	1020	1030	1040	
OnInR	YEISVQACRKTEPDKGD-----FQPNCSDSTSKEIRTESKHDADNIASNISIVSVTEN----SITISWK-PPINPNGRI								871
DInR	YAIFVACREEIPSEK--LRDTSFKKSLCSDYDTVFQTTKRKKFADIVMDLKVDLEHANNTESTPATFSLGRHQLDPNGEI								1235
HIR	YRIELQACNQDTP-----EERCSSVAAYVSARTMPEAKADDIVGPVTHEIFENN----VVHLMWQ-EPKEPNGLI								881
AaeInR	YVFKIRACRHPGDPAPSVRLVDVELACGNEVFENFRTPKKEGADDIPPESILIEEQSNNTQRQIRVQWK-EPKPNGPI								869
BomInR	YTVNIWACRNKDLNETDPV---YAENWCSVRAYNTRTLELPPNAD-VVSNFQVEIMPANKTLPEVNVNWE-PPLNPNGV								941
Majority	VXVEVXYSRXGDEXV-----	LXXCIXXXDFNQTR-GYLLKXLSPGNYSXRVRAXSIAGXGXXEAXYVXVXXXDEXSXX							
		+-----+-----+-----+-----+-----+-----+-----+							
	1050	1060	1070	1080	1090	1100	1110	1120	
OnInR	RHYFIDVSKSGTPIP-----ERCIDNQDFKNSSYKYTIEPLSPGNYTIRVQAISSANTG---GAFSLPVSYYIDEPSDS								942
DInR	VTYEVAYKLQKPDQVE----EKKCIPAAFNFQTA-GYLK-LNEGLYFRVRANSIAGYGDFTEVEHIKVEP-PPSYAKV								1308
HIR	VLYEVSYRRYGDEE-----LHLCVSRKHAFALER-GCRLRGLSPGNYSVRIRATSLAGNGSWTEPTYFYVTDYLDVPSNI								954
AaeInR	VKFVVKYQRVDLESVSS----TDICIRYSSFNQTR-GALLTKLEPGNYSIRVMATTIAGDGAPSAARYVLIAM--DDSMGT								943
BomInR	VAYNVHYSRIEDSAQGPVGLQSCITVDDYEANGHGYLTKLSPGNYSVKVTPITVSGAGNVSEEIYVFIPEPVSESGHA								1021

Supplementary Figure 7. Insulin receptor protein alignment of the Fibronectin type III domains. Abbreviations: *O. niriventrtris* insulin receptor (OnInR), *Drosophila* insulin receptor (DInR), human insulin receptor (HIR), *A. aegypti* insulin receptor (AaeInR), *B. mori* insulin receptor (Bom InR).

Majority	WXIILG--GLXXLFLXVVGXVXYXLRKKRPPXEXRMXXSVNPEYAS-----XYVPDDWEVXREKIIXLRELQGGSF	
OnInR	WNIIAG---IIILVLIIVGIIIVYILFKKRQEEESPRMNP SINPEYM-----YVPDDWEIPRKKIELHKELQGGSF	1009
DInR	FFWLLG-IGLAFILVSLFGYVCYLHKKRVPSND-LHMNTEVNP FYAS-----MQYIPDDWEVLRENIIQLAPLQGGSF	1379
HIR	AKIIIIGPLIFVFLFSVVIGSIYLFLRKRQPDGPLGPLYASSNPEYLSASDVFPCSVYVPDEWEVSREKITLLRELQGGSF	1034
AaeInR	TLIWLG--TLIVIFLCSVGFVAFYWKYRYMSKQIRMPYEVNPDYAG-----VQYKVDDEVERNHIIQLEELQGGSF	1014
BomInR	WAWGVG-AGGVVLALLVGAADVYARRGLMSHAEASKL FASVNPEYVPT-----VYVPDEWEVTRDSIHFIRELQGGSF	1093
Majority	GMVYEGIXDXIXKGAETRCAIKTVNEXATDRERIXFLNEASVMKAFDTHHVRL LGVVSXGQPTLVVMELMANGDLKSY	
OnInR	GMVYEGIVRDIK-GKATIRCAIKTVNEHATNAERLNFLKEASVMKAFDTHAVIKLLGVVSXGQPTLVVMELMANGDLKSY	1088
DInR	GMVYEGILKSFPPNGVDRECAIKTVNENATDRERTNFLSEASVMKEFDTYHVVRL LGVCSRGPALVVMELMCKGDLKSY	1459
HIR	GMVYEGNARDIIKGEAETRVAVKTVNESASLRERIEFLNEASVMKGFTHHVRL LGVSKGQPTLVVMELMAHGDLSY	1114
AaeInR	GMVYKGI L TQLRGEKCNQPCA IKT VNESATAREKDSFLLEASVMKQFNTHHVRL LGVVSQGDPTLVIMELMANGDLKSY	1094
BomInR	GMVYEGIAKEIVKGPETRCAIKTVNEHATDRERIEFLNEASVMKAFDTHHVRL LGVVSXGQPTLVVMELMERGDLSY	1173
Majority	LRSRHPDAEG-----NXPGSPX--QPPTLXRIXQMAIEIADGMAYLSAKKFVHRDLAARNCMVADDLTVKIGDFGMTR	
OnInR	LRLHRPDS DG-----YTHGIIG--QPPSLRRILQMAIEIADGMAYLSAKKFVHRDLAARNCMVAEDLTVKIGDFGMTR	1159
DInR	LRAHRPEERDEAMMTYLNRI GVTGNVQPPTYGRIYQMAIEIADGMAYLAAKKFVHRDLAARNCMVADDLTVKIGDFGMTR	1539
HIR	LRLSRPEAE-----NNPGRP----PPTLQEMIQMAAEIADGMAYLNAKKFVHRDLAARNCMVAHDFTVKIGDFGMTR	1182
AaeInR	LRRHRPDYEN-----GEDPSP---QPPTLRQIIQMAIEIADGMAYLSAKKFVHRDLAARNCMVADDMTVKIGDFGMTR	1164
BomInR	LRSRHPDAEG-----SLPSSPP--VPPTLQNILQMAIEIADGMAYLSAKKFVHRDLAARNCMVAGDLTVKIGDFGMTR	1244
Majority	DIYETDYRKGTKGLLPVRWMAPESLKDGVSFSSSDVSYGVVLEWEMATLASQPYQGLSNEQVLRVYVIDGGVMERPENC	
OnInR	DIYETDYRKGTKGLLPVRWMAPESLKDGVSFSSSDIWSYGVVLEWEMATLASQPYQGLSNDQVLRVYVIDSGIMERPENC	1239
DInR	DIYETDYRKGTKGLLPVRWMPPEL RDGVYSSASDVFSFGVLEWEMATLAAQPYQGLSNEQVLRVYVIDGGVMERPENC	1619
HIR	DIYETDYRKGTKGLLPVRWMAPESLKDGVFTTSSDMWSFGVLEWETS LAEQPYQGLSNEQVLFVMDGGYLDQPDNCP	1262
AaeInR	DIYETDYRKGTKGFLPVRWMAPESLKDGIFSSSDVFSYGVVLEWEMATLASQPYQGLTNDQVLRVYVIDGGVMERPENC	1244
BomInR	DIYETDYRKGTKGLMPVRWMSPEL KDGVSFSSSDAWSYGVVLEWEMATLAMAQPYQGLSNEQVLRVYVVEGGVMERPEQCP	1324
Majority	DRLYXLMRRCWQHRSARPTF-LXIVXXLLPDA-SPQFXEVSFYHSEAG-----XRAXXREE-----Q	
OnInR	DKLYHLMRQCWKYKYAERPSFXLDLVSMLLEDS-SPQFSRVSFYHSAAG-----MEARSTRN-----N	1296
DInR	DFLHKL MQR CWHRSSARPSF-LDIIAYLEPQC PNSQFKEVSFYHSEAG-----LQHREKERKERNQLDAFAAVPLDQDLQ	1694
HIR	ERVTDLMRMCWQFNPKMRPTF-LEIVNLLKDDL-HPSFPEVSFFHSEE-----NKAPESEE-----L	1317
AaeInR	DNLYNLMRRCWQHRTARPTF-MEIISELLPDA-SPHFQDVAFYNSQD-----ALDMLRGQ-----H	1299
BomInR	DRLYELMRACWTHRPSARPTF-LQLVADLVPHA-MPYFRHRSFFHSPQGGELYALQRSALDEE-----Q	1386
Majority	XMEIQDDATXPLR-----	
OnInR	QMIIQDDVMMPLR-----	1309
DInR	DREQQEDATPLRMGDYQQNSSL DQPPEPIAMVDDQGSHPFSLPSGFIASSTPDGQTVMATAFQNI PAAQGDISATYV	1774
HIR	EMEFEDMENVPLD-----	1330
AaeInR	QTVIIDEATTPLR-----	1312
BomInR	ELPEVDVGAVATGS-----	1400

Majority	-----XXSXXXREXPGLRSGGS-----XSGXXDDLXGGPPXEFS-AYTXSXX	
	-----+-----+-----+-----+-----+-----+-----+	
OnInR	-----LTKEL E E I Y P L N S C A D-----DSEVEDDLVGSPHMEFS-SYPKVQK	1349
DInR	VPDADALDGRGYE I Y D P S P K C A E L P T S R S G S T G G G K L S G E Q H L L P R K G R Q P T I M S S S M P D D V I G G S S L Q P S T A F T T S S N	1854
HIR	-----RSSH C Q R E E A G G R D G G S-----S L G F K R S Y E E H I P-----Y T H M N G	1366
AaeInR	-----P G D D H D E E P G E D D D L V-----G H G E G H I G D V G T D D E F S M E M T N S H L	1353
BomInR	-----V S N L F G V S G R L A S W R E L S S L R S R G S-----D D A A A E P L Q P L P P P G T S P A P I L A P A	1451
Majority	XXXNGPXXTLRXXXXSXXXX-----	
	-----+-----+-----+-----+	
OnInR	DTTGTANGYVGTNPSNGTRA	1370
DInR	ASSHTGRPSLKKTVADSVRNKANFINRHLY	1884
HIR	GKKNRILTLPRSNPS	1382
AaeInR	VRNNGPMATIRSPHSPLR	1371
BomInR	PAIKGPNGLRDSHSARFRNH	147

Supplementary Figure 8. Insulin receptor protein alignment of the transmembrane, juxtamembrane, and tyrosine kinase domains. Abbreviations: *O. nigriventris* insulin receptor (OnInR), *Drosophila* insulin receptor (DInR), human insulin receptor (HIR), *A. aegypti* insulin receptor (AaeInR), *B. mori* insulin receptor (BomInR).



Supplementary Figure 9. Distribution of OnInR in late prepupal stage imaginal tissues. Lanes 2-5: large male (LM) horn (H), wing (W), leg (L), and genital (G) imaginal discs. Lanes 6-9: small male (SM) H, W, L, G. Lanes 10-12: large female (LF) H, W, L. Lanes 13-15: small female (SF) H, W, L. Molecular weight markers (LD) in nucleotides are indicated.